

Strong Haken via Sphere Complexes

Sebastian Hensel and Jennifer Schultens

February 1, 2023

Abstract

We give a short proof of Scharlemann’s Strong Haken Theorem for closed 3-manifolds (and manifolds with spherical boundary). As an application, we also show that given a decomposing sphere R for a 3-manifold M that splits off an $S^2 \times S^1$ summand, any Heegaard splitting of M restricts to the standard Heegaard splitting on the summand.

1 Introduction

Any (closed, oriented, connected) three-dimensional manifold M admits a *Heegaard splitting*, that is, it can be decomposed into two three-dimensional handlebodies V, V' of the same genus g along an embedded surface $S \subset M$:

$$M = V \cup_S V'.$$

In theory, all information about the three-manifold is encoded in the identification of the two handlebodies. However, in practice, interpreting topological properties of M using a Heegaard splitting is often nontrivial.

A basic example of this occurs when studying spheres in M . If $\alpha \subset S$ is a curve which bounds disks D, D' in both V and V' , then gluing these disks yields a 2-sphere $D \cup D' \subset M$ which intersects S in the single curve α . When essential, such a sphere is called a *Haken sphere* – but a priori it is completely unclear what kind of spheres in M are of this form.

A classical theorem of Haken [5] shows that if M admits any essential sphere σ , then it also admits a Haken sphere σ' . In fact, Scharlemann [16] recently proved a *Strong Haken Theorem*, showing that σ' can in fact be chosen to be isotopic to σ :

Theorem 1.1. (*Strong Haken Theorem*) *Let $M = V \cup_S V'$ be a Heegaard splitting. Every essential 2-sphere in M is isotopic to a Haken sphere for $M = V \cup_S V'$.*

The purpose of this article is to give an independent, short proof of Theorem 1.1 for any M which is closed or has spherical boundary. We want to mention that Scharlemann’s version of the strong Haken theorem is in fact more general, allowing for manifolds with arbitrary boundary, and also showing that any properly embedded disk is isotopic to a Haken disk. This more general case could also be obtained from our methods; for clarity we focus on the closed case throughout the article.

To prove Theorem 1.1, we make crucial use of the (*surviving*) *sphere complex*, which is a combinatorial complex encoding the intersection pattern of essential (surviving) spheres

in M . Such complexes have already been used successfully in the study of outer automorphism groups of free groups (via mapping class groups of connected sums of $S^2 \times S^1$). Here, we show that this perspective can also be useful in streamlining arguments in low-dimensional topology. The other crucial ingredient is the classical Waldhausen theorem on Heegaard splittings of the three-sphere [19]. Together, these allow an inductive approach to Theorem 1.1.

Our methods and results also allow to control Heegaard splittings of reducible manifolds. As a motivating example, we prove:

Proposition 1.2. *Every Heegaard splitting of $W_n = n(S^2 \times S^1)$ is isotopic to a stabilization of the standard Heegaard splitting.*

Combining the uniqueness for W_1 with the Strong Haken Theorem, we obtain the following structural result on Heegaard splittings of arbitrary reducible three-manifolds.

Corollary 1.3. *Given a reducible 3-manifold M with a Heegaard splitting $M = V \cup_S V'$, any decomposing sphere that splits off a $S^2 \times S^1$ summand can be isotoped so that S is standard in this summand.*

Acknowledgements We would like to thank Martin Scharlemann for finding a mistake in an earlier draft, and many helpful comments.

2 Heegaard Splittings of Closed 3–Manifolds

In this section we recall some preliminaries on closed three-manifolds, their Heegaard splittings, and spheres in such manifolds. The results presented here are classical.

2.1 Heegaard Splittings

Definition 2.1. (*Heegaard splitting*) *A handlebody is a 3-manifold that is homeomorphic to a regular neighborhood of a graph in S^3 . A Heegaard splitting of a 3-manifold M is a decomposition $M = V \cup_S V'$, where V, V' are handlebodies and $S = \partial V = \partial V' = V \cap V'$.*

The surface S is called the splitting surface or Heegaard surface. Heegaard splittings are considered equivalent if their splitting surfaces are isotopic.

Remark 2.2. *The Heegaard splitting $M = V \cup_S V'$ is completely specified by the pair (M, S) , so we will sometimes write (M, S) instead of $M = V \cup_S V'$.*

Remark 2.3. *The connected sum of two 3-manifolds M_1, M_2 with Heegaard splittings $(M_1, S_1), (M_2, S_2)$ inherits a Heegaard splitting $(M_1 \# M_2, S_1 \# S_2)$. This Heegaard splitting is unique in the sense that it is completely determined by the construction. Later, we will briefly consider a refined notion of equivalence for Heegaard splittings where we distinguish between the two sides of the splitting surface. With respect to this refined notion of equivalence, the Heegaard splitting of a connected sum is then not (a priori) unique, as the construction allows two different choices, namely which side of S_1 is identified to which side of S_2 .*

Definition 2.4. Given a Heegaard splitting (M, S) , the Heegaard splitting obtained from the pairwise connected sum $(M, S) \# (S^3, T)$, where T is the standard unknotted torus in S^3 , is called a stabilization of (M, S) . A Heegaard splitting is stabilized if it is the stabilization of another Heegaard splitting and unstabilized otherwise.

A sphere that separates $(M, S) \# (S^3, T)$, i.e., a sphere that splits off a punctured 3-ball containing an unknotted punctured torus, is called a stabilizing sphere.

A stabilizing pair of disks is a pair (D, D') of disks such that D is properly embedded in V , D' is properly embedded in V' and $\partial D \cap \partial D'$ is exactly one point.

Remark 2.5. A Heegaard splitting is stabilized if and only if it admits a stabilizing pair of disks. Indeed, consider the standard unknotted torus in the 3-sphere and observe that it separates S^3 into two solid tori. The boundaries of the meridian disks of these solid tori intersect in exactly one point.

A crucial theorem of Waldhausen's characterises all Heegaard splittings of the three-sphere. See [19].

Theorem 2.6 (Waldhausen's Theorem). *Every Heegaard splitting of the three-sphere is a stabilization of the unique standard genus 0 Heegaard splitting.*

2.2 Sphere Complexes

A core tool in our argument is the following simplicial complex, which encodes the intersection patterns of spheres in M .

Definition 2.7 (Sphere complex). *A sphere S in a 3-manifold is compressible if it bounds a 3-ball. Otherwise, it is incompressible. We say that a sphere is peripheral if it is isotopic into the boundary of the manifold.*

The sphere complex of a 3-manifold M is the simplicial complex $\mathcal{S}(M)$ determined by the following three conditions:

1. Vertices of $\mathcal{S}(M)$ correspond to isotopy classes of incompressible, non-peripheral embedded 2-spheres;
2. Edges of $\mathcal{S}(M)$ correspond to pairs of vertices with disjoint representatives;
3. The complex $\mathcal{S}(M)$ is flag.

It is not hard to see that a simplex in the sphere complex corresponds to a collection of nonisotopic spheres that can be realised disjointly. Compare Figure 1 for an example of a simplex.

Furthermore, a standard argument involving surgery at innermost intersection circles shows that the sphere complex of any closed 3-manifold is connected (if it is nonempty). See e.g., [6] for a proof in the case of doubled handlebodies, which also works in general.

2.3 Haken Spheres

Our central aim will be to understand how essential spheres in M interact with Heegaard splittings of M . The following notion is crucial.




Figure 1: A 3-simplex in the sphere complex of W_4 , the double of a genus 4 handlebody (alternatively, the connected sum of 4 copies of $S^2 \times S^1$). Here, only one of the handlebodies is pictured; the spheres comprising the simplex intersect it in the pictured disks.

Definition 2.8. Let $M = V \cup_S V'$ be a Heegaard splitting. An essential sphere in M that meets the Heegaard surface S in a single simple closed curve is called a Haken sphere. A (not necessarily essential) sphere that intersects S in a single simple closed curve essential in S is called a reducing sphere.

The following theorem was originally proved by Haken in [5]. Proofs can be found in the standard references on 3-manifolds, see [7], [8], [18].

Theorem 2.9 (Haken's Lemma). *If a closed three-manifold M contains an essential sphere and $M = V \cup_S V'$ is a Heegaard splitting, then M admits a Haken sphere.*

In general, the Haken sphere is obtained by modifying the given essential sphere by surgery, and so cannot be guaranteed to be related to the sphere given at the outset.

3 Three-manifolds with spherical boundary

In this section, we present versions of the results and notions of the previous section for 3-manifolds with spherical boundary. These appear naturally in our inductive proof of the Strong Haken Theorem (even if one is just interested in proving it in the closed case). For ease of notation, if M is a 3-manifold with spherical boundary, then we call each component of ∂M a *puncture*. Similarly, we call such a manifold a *punctured manifold*.

3.1 Heegaard Splittings

To define Heegaard splittings of punctured manifolds, we use spotted handlebodies.

Definition 3.1. A spotted handlebody is a handlebody with a specified set of disks $D_1 \sqcup \cdots \sqcup D_k$ in its boundary. Each disk is called a spot. A Heegaard splitting of a 3-manifold M with spherical boundary is a decomposition $M = V \cup_S V'$ where V, V' are spotted handlebodies with spots $D_1 \sqcup \cdots \sqcup D_k$ and $D'_1 \sqcup \cdots \sqcup D'_k$, respectively, and $S = \partial V - (D_1 \sqcup \cdots \sqcup D_k) = \partial V' - (D'_1 \sqcup \cdots \sqcup D'_k)$.

Remark 3.2. *In a Heegaard splitting of a 3-manifold with spherical boundary each puncture meets the splitting surface in a single simple closed curve. This simple closed curve is the boundary of a spot on each of the handlebodies.*

Suppose M_1, M_2 are two punctured manifolds, with boundary components $\partial_i \subset M_i$, and $M = M_1 \cup_{\partial_1 = \partial_2} M_2$ is the manifold obtained by gluing the boundary components. Given Heegaard splittings of M_1, M_2 , the manifold M inherits a Heegaard splitting which is obtained by gluing the handlebodies at the corresponding spots.

The glued boundary components yield an essential 2-sphere σ in M , which intersects the induced Heegaard splitting in a single circle (i.e., it becomes a Haken sphere). Conversely, given a Haken sphere σ for any manifold M , one can cut the manifold and the splitting at σ .

We need a version of Waldhausen's Theorem in the context of punctured 3-spheres (which is a fairly straightforward consequence of Waldhausen's theorem for S^3).

Theorem 3.3 (Waldhausen's Theorem for punctured 3-spheres). *Every Heegaard splitting of a punctured 3-sphere is a stabilization of a unique standard genus 0 Heegaard splitting.*

Proof. Let M be a punctured 3-sphere and $M = V \cup_S V'$ a Heegaard splitting. Construct $\hat{M} = S^3$ from M by attaching a 3-ball to each puncture. By Alexander's Theorem, the result does not depend on the attaching map. Moreover, the attaching maps can be chosen so that a meridional disk of each 3-ball caps off a component of ∂S . We thus obtain a closed surface \hat{S} that defines a Heegaard splitting $S^3 = \hat{V} \cup_{\hat{S}} \hat{V}'$.

Each 3-ball that has been attached to a puncture is a regular neighborhood of a point and, as such, arbitrarily small. By Waldhausen's Theorem, $S^3 = \hat{V} \cup_{\hat{S}} \hat{V}'$ is a stabilization of the standard genus 0 Heegaard splitting of S^3 . The stabilizing pairs of disks can be chosen to be disjoint from the attached 3-balls. Thus, after destabilizing, if necessary, we may assume that S is genus 0.

Hence, to prove the theorem, it suffices to show that any genus 0 splitting of a punctured S^3 is standard. To this end, observe that the spotted genus 0 handlebody $V \subset S^3$ can be isotoped to be a regular neighbourhood of a graph $\Gamma \subset V$. The graph Γ can be chosen to have the following form: it has one vertex v_0 in the interior of M , and one vertex on each boundary component. Each vertex on a boundary component is joined to v_0 by an edge. Now, observe that any two such graphs are isotopic, as any two arcs from v_0 to a boundary sphere are isotopic, by the Lightbulb Trick. This shows that any two genus 0 splittings of a punctured S^3 are isotopic. \square

At this point, we briefly want to address the ambiguity appearing in the previous proof when filling the boundary and drilling it out again – namely, one can isotope a pair of stabilizing disks across a puncture. This leads to a homeomorphism of the manifold preserving the Heegaard surface. Given a Heegaard splitting of a 3-manifold, the *Goeritz group* of the splitting is the group of isotopy classes of orientation preserving diffeomorphisms of the manifold that preserve the splitting. Loosely speaking, the Goeritz group will be small if the surface automorphism that defines the Heegaard splitting is complicated relative to the handlebodies. Conversely, the Goeritz group will be as large as possible in the case of W_n , the manifold for which this surface automorphism is the identity, and the Goeritz group is equal to the handlebody group. Scharlemann finds a system of $4g + 1$ generators for the Goeritz group of a handlebody (see [15]). On the other hand, the Goeritz group of the

three-sphere is still largely mysterious. We refer the interested reader to the recent [17] and the references therein.

3.2 Sphere Complexes

We now want to define a useful sphere complex for punctured manifolds. One obvious change is that for the vertices one should also exclude *peripheral* spheres, i.e., spheres which are homotopic into the boundary (otherwise, such spheres are adjacent to any other vertex, rendering the resulting complex useless). However, even with this modification, the resulting sphere complex will be somewhat problematic for our purposes, as it may often be disconnected. Namely, suppose that M_0 is an aspherical three-manifold with infinite fundamental group. Let M be the manifold obtained from M_0 by removing two open balls. The manifold M admits many essential non peripheral spheres obtained by joining the two punctures by a nontrivial tube. In fact, by asphericity of M_0 , any essential non peripheral sphere in M is of this form. In particular, no two such are disjoint.

To sidestep this issue, we use the following variant of the sphere complex.

Definition 3.4 (Surviving Sphere complex). *A sphere S in a punctured 3-manifold is almost peripheral if it bounds a punctured S^3 in M . If S is not almost peripheral, then it is surviving.*

The surviving sphere complex of a 3-manifold M is the simplicial complex $\mathcal{S}^s(M)$ determined by the following three conditions:

1. *Vertices of $\mathcal{S}^s(M)$ correspond to isotopy classes of incompressible, embedded, surviving 2-spheres;*
2. *Edges of $\mathcal{S}^s(M)$ correspond to pairs of vertices with disjoint representatives;*
3. *The complex $\mathcal{S}^s(M)$ is flag.*

The terminology stems from the fact that the spheres “survive filling in the punctures” and is in analogy to the surviving curve complex used in the study of mapping class groups of surfaces, see e.g., [1, 4]. It turns out that these complexes are much better behaved in our setting.

Lemma 3.5. *Let M be a 3-manifold. Then the surviving sphere complex $\mathcal{S}^s(M)$ is connected (if it is nonempty).*

Proof. Let σ, σ' be two incompressible, embedded, surviving 2-spheres in M . Up to isotopy, we may assume that σ, σ' intersect transversely. Further, we may assume that up to isotopy, the number of intersection components $\sigma \cap \sigma'$ is minimal.

Let $C \subset \sigma \cap \sigma'$ be an innermost intersection circle, i.e., suppose that it bounds a disk $D \subset \sigma$ with $D \cap \sigma' = \partial D = C$. Denote by $S^+, S^- \subset \sigma'$ the two disks bounded by C , and denote by $\sigma^\pm = S^\pm \cup D$ the two 2-spheres obtained by disk-swapping. Observe that up to isotopy, both of these are disjoint from σ' , and intersect σ in at least one fewer circle than σ' . If either σ^+, σ^- were compressible, then we could reduce the number of components in $\sigma \cap \sigma'$ by sliding σ over the ball bounded by the compressible sphere, which is impossible by our choice.




Figure 2: In the proof of Lemma 3.5: the innermost intersection circle of σ, σ' cuts σ' into two disks S^+, S^- . If σ^- is almost peripheral, then σ^+ is isotopic to σ after filling the punctures.

Assume that σ^- is almost peripheral. Then, after filling in the punctures of M , the spheres σ and σ^+ are isotopic (by sliding D over the now unpunctured ball bounded by σ^- ; compare Figure 2). In particular, σ^+ is surviving, as the same is true for σ .

Hence, at least one of σ^\pm is surviving, and we are done. Indeed, repeating this process produces a sequence of spheres corresponding to vertices in a path, in $\mathcal{S}^s(M)$, between $[\sigma]$ and $[\sigma']$. \square

3.3 Haken Spheres

Just as in the closed case, we call an essential sphere which intersects a Heegaard splitting of a punctured manifold in a single curve a *Haken sphere*. For punctured manifolds, almost peripheral and surviving Haken spheres behave slightly differently.

On the one hand, using the same strategy as in the proof of Theorem 3.3, we obtain the following corollary of Theorem 2.9.

Theorem 3.6 (Surviving Haken's Lemma). *If M contains a surviving sphere and $M = V \cup_S V'$ is a Heegaard splitting, then there is a surviving Haken sphere.*

Proof. Denote by σ an essential surviving sphere in M . Let M' be the three-manifold obtained from M by gluing a ball to each boundary component. Denote by $B \subset M'$ the disjoint union of the resulting balls. By definition of almost peripheral, the image of σ in M' is still essential. Thus, Haken's lemma (Theorem 2.9) applies, and yields a Haken sphere $\sigma' \subset M'$. By an isotopy preserving the Heegaard surface, we may assume that σ' is disjoint from B . We can thus interpret σ' as a sphere in $M \subset M'$, where it is the desired Haken sphere. \square

On the other hand, almost peripheral spheres are also Haken spheres:

Lemma 3.7 (Almost Peripheral Strong Haken Theorem). *Let M be a three-manifold with at least two punctures, and $M = V \cup_S V'$ be a Heegaard splitting. Then any almost peripheral sphere σ in M is isotopic to a Haken sphere.*

Proof. We begin with the case where σ cuts off exactly two punctures δ_1, δ_2 . The almost peripheral sphere σ is then isotopic to the boundary of a regular neighbourhood of $\delta_1 \cup \alpha \cup \delta_2$, where $\alpha \subset M$ is a properly embedded arc. We may homotope α to lie in S , as any arc in a




Figure 3: *Removing self-intersections of an arc joining two spots.*

handlebody is homotopic into the boundary. However, the arc may now not be embedded anymore. We can remove the self-intersections by “popping subarcs over $\delta_1 \cap S$ ”. To be more precise, parametrise $\alpha : [0, 1] \rightarrow S$ so that it starts on $\delta_1 \cap S$, and homotope so that all self-intersections are transverse. Consider the first self-intersection point $\alpha(t) = \alpha(s), t < s$. In particular, this implies that $\alpha|_{[0,t]}$ is an embedded arc.

Now homotope a small subarc $\alpha|_{[s-\epsilon, s+\epsilon]}$ to instead be the arc obtained by following $\alpha|_{[0,t]}$ backwards to $\delta_1 \cap S$, following around $\delta_1 \cap S$, and returning along $\alpha|_{[0,t]}$ (compare Figure 3). This homotopy is possible in V , and the resulting arc has at least one fewer self-intersection.

After a finite number of modifications of this type, the boundary of a regular neighbourhood of $\delta_1 \cup \alpha \cup \delta_2$ (which is homotopic to σ) intersects S in a single curve, and thus is a Haken sphere. By a theorem of Laudenbach, see [9], homotopy and isotopy are the same for spheres in 3-manifolds, hence the claim follows¹.

Now we suppose σ is a sphere cutting off $k > 2$ punctures. Then there is a sphere σ' , disjoint from σ , which cuts off 2 punctures, and which is contained in the punctured S^3 bounded by σ . By the initial case, σ' may be assumed to be Haken. Let M' be the manifold obtained by cutting M at σ' , with the induced Heegaard splitting; observe that $\sigma \subset M'$ is still almost peripheral, but now cuts off at most $k - 1$ spheres. By induction, σ is a Haken sphere. \square

Since any essential, non peripheral sphere in a punctured S^3 is almost peripheral, this implies the following:

Corollary 3.8 (Strong Haken Theorem for punctured 3-spheres). *Any essential sphere in a punctured 3-sphere is isotopic to a Haken sphere.*

4 Heegaard splittings of $n(S^1 \times S^2)$

In this section, we study Heegaard splittings of a specific manifold, namely

Definition 4.1. *We denote the double of the genus n handlebody by W_n . It is the connected sum of n copies of $S^2 \times S^1$.*

¹One could also avoid citing this theorem by isotoping α into a regular neighbourhood of S and resolving crossings of the projection to S by isotopies which slide strands over the puncture similar to Figure 3.

A reader only interested in the strong Haken Theorem may safely skip ahead to the next section. Our goal here will be to prove that, similar to Waldhausen’s Theorem for the three-sphere, all Heegaard splittings of W_n are “standard” in the following sense.

Definition 4.2. *A Heegaard splitting of W_n is standard if it is the double of a genus n handlebody. A standard Heegaard splitting of W_n is a Heegaard splitting that is the connected sum of n copies of W_1 with the standard Heegaard splitting.*

Waldhausen seems to claim in [19] that all Heegaard splittings of W_n are standard (although it is not entirely clear up to which equivalence relation, and the proof sketch is incomplete). In the unpublished preprint [12], Oertel and Navarro Carvalho prove the result, using results on the homeomorphism groups of handlebodies and W_n (in a very similar way to the argument we will use below). In this section, we show that these techniques could also be used to prove a Strong Haken Theorem (and obtain the uniqueness of splittings as a corollary). We want to emphasize that this of course follows from the general Strong Haken Theorem (Theorem 1.1), but consider the argument using homeomorphisms of W_n interesting enough to warrant this alternate proof.

Proposition 4.3. *Every unstabilized Heegaard splitting of W_1 is standard.*

Proof. Suppose that $W_1 = V \cup_S V'$ is a Heegaard splitting. We wish to show that $W_1 = V \cup_S V'$ is standard. Since W_1 is reducible, Haken’s lemma tells us that there is a Haken sphere R for $W_1 = V \cup_S V'$. Denote $V \cap R$ by D and $V' \cap R$ by D' . Note that all essential spheres in W_1 , in particular R , are isotopic to $S^2 \times (\text{point})$.

We may assume that S intersects a bicollar of R in an annulus $(S \cap R) \times [-1, 1]$. Removing this bicollar leaves a copy of $S^2 \times [-1, 1]$, i.e., a twice punctured 3-sphere that inherits a Heegaard splitting. By Theorem 3.3, this Heegaard splitting is either of genus 0 or stabilized.

Since $W_1 = V \cup_S V'$ is unstabilized, the Heegaard splitting obtained on the complement of $S^2 \times [-1, 1]$ must be of genus 0. Specifically, the splitting surface is a twice punctured 2-sphere, i.e., an annulus. Hence we can reconstruct $W_1 = V \cup_S V'$: Indeed, say, V , is composed of a 3-ball attached to the two copies $D \times \{\pm 1\}$ of D . It follows that V is a solid torus. The same is true of V' and hence $W_1 = V \cup_S V'$ is the standard Heegaard splitting. \square

First, we have the following classical result due to Griffiths [3].

Theorem 4.4. *The action of the mapping class group of a handlebody V_n on its fundamental group $\pi_1(V_n) = F_n$ induces a surjection*

$$\text{Mcg}(V_n) \rightarrow \text{Out}(F_n) \rightarrow 1.$$

We remark that the kernel of this map is quite complicated, and generated by twists about disk-bounding curves [11]. Next, we need a theorem of François Laudenbach [9] (see also [10] and [2] for a modern proof):

Theorem 4.5. *The action of the mapping class group of a doubled handlebody W_n on its fundamental group $\pi_1(W_n) = F_n$ induces a short exact sequence*

$$1 \rightarrow K \rightarrow \text{Mcg}(W_n) \rightarrow \text{Out}(F_n) \rightarrow 1.$$

The kernel K is finite, generated by Dehn twists about non-separating spheres, and acts trivially on the isotopy class of every embedded sphere or loop.

Corollary 4.6. *For the standard Heegaard splitting of W_n , every essential sphere in W_n is isotopic to a Haken sphere.*

Proof. First observe that any two nonseparating spheres in W_n can be mapped to each other by a homeomorphism. Namely, the complement of such a sphere is homeomorphic to W_{n-1} with two punctures. Similarly, separating spheres can be mapped to each other if and only if the fundamental groups of the complements are free groups of the same rank (as the complement is a disjoint union of once-punctured W_k and W_{n-k}).

Next, observe that there are Haken spheres of all such possible types, obtained by doubling a suitable disk in the handlebody.

Let $i : V_n \rightarrow W_n$ be the inclusion induced by doubling. Observe that on the one hand, the boundary of V_n maps under i to the standard Heegaard splitting of W_n , and on the other hand i induces an isomorphism i_* of fundamental groups. For any outer automorphism $\varphi \in \text{Out}(\pi_1(V_n))$ of the fundamental group of V_n , by Theorem 4.4, there is a homeomorphism $f : V_n \rightarrow V_n$ inducing it. Let $F : W_n \rightarrow W_n$ be the homeomorphism of W_n obtained by doubling f . Observe that F preserves the standard Heegaard splitting of W_n by construction, and F induces φ via the isomorphism $i_* : \pi_1(V_n) \rightarrow \pi_1(W_n)$. Since φ was arbitrary, this shows that any outer automorphism of $\pi_1(W_n)$ can in fact be realised by a homeomorphism of W_n preserving the standard Heegaard splitting.

Together with Laudenbach's Theorem 4.5 this shows that any sphere is isotopic to the image of a Haken sphere under a homeomorphism preserving the standard Heegaard splitting – hence, it is isotopic to a Haken sphere. \square

Lemma 4.7. *There is a unique Heegaard splitting of W_n of genus n .*

Proof. Connected sum decompositions of W_n are not unique. However, let $W_n = V \cup_S V'$ be the standard Heegaard splitting and let $W_n = X \cup_Y X'$ be any Heegaard splitting of genus n . By repeated application of Theorem 2.9 there are Haken spheres $R_1 \cup \dots \cup R_{n-1}$ for W_n that cut $W_n = X \cup_Y X'$ into standard Heegaard splittings of W_1 . By Corollary 4.6, R_1, \dots, R_{n-1} are also Haken spheres for $W_n = V \cup_S V'$. By an Euler characteristic argument, these cut $W_n = V \cup_S V'$ into genus 1 Heegaard splittings of the summands. By Proposition 4.3 these are standard. In particular, Y is isotopic to S . \square

Proof of Proposition 1.2. For an unstabilized Heegaard splitting, this is Lemma 4.7. Furthermore, if $n = 1$, then this follows from Proposition 4.3. So suppose that $n > 1$ and $W_n = V \cup_S V'$ is stabilized. By Corollary 4.6, there is a Haken sphere R that decomposes W_n into $W_1 \# W_{n-1}$. Moreover, by [13], one of the Heegaard splittings inherited by the summands is stabilized. By induction, $W_n = V \cup_S V'$ is a stabilization of a connected sum of standard Heegaard splittings, i.e., a stabilization of the standard Heegaard splitting of W_n . \square

5 Strong Haken Theorem

Combining the uniqueness of Heegaard splittings for W_n (Proposition 1.2) with Corollary 4.6 yields a *Strong Haken Theorem* for W_n : any sphere in W_n is isotopic to a Haken sphere. This statement was recently proved by Scharlemann [16] for all three-manifolds. In this

section, we provide a short independent proof of this theorem for closed manifolds and manifold with spherical boundary.

The following proof proceeds by two nested inductions. It naturally involves 3-manifolds with spherical boundary, even if we just want to prove the theorem in the closed case. Recall that for such 3-manifolds, we decree that each boundary sphere (puncture) meets the splitting surface in a single simple closed curve.

Proof of Theorem 1.1. We prove the theorem by considering all punctured 3-manifolds and all Heegaard splittings, ordered according to a suitable complexity. Namely, if $M = V \cup_S V'$ is a Heegaard splitting, we define the *complexity* as the pair $(g(S), n(S))$ of genus and number of spots of the handlebodies (ordered lexicographically). We perform a nested induction on the genus g and the number of boundary components n . The argument for the inductive step is in fact the same in both cases, and so we describe both inductions simultaneously.

Induction Start $g = 0, n \geq 0$ Observe that the only punctured 3-manifold that can be obtained from a Heegaard splitting of genus 0 is the three-sphere. Thus, the Strong Haken Theorem in this case is simply Corollary 3.8.

Induction Steps Now suppose that the Strong Haken Theorem is known for all manifolds of complexity at most (g, n) , and suppose that M is a manifold of complexity $(g, n + 1)$, or suppose that the Strong Haken Theorem is known for all manifolds of complexity $(g, k), k \geq 0$, and M is a manifold of complexity $(g + 1, k)$.

First observe that, by Lemma 3.7 any almost peripheral sphere in M is isotopic to a Haken sphere. We thus have to show that surviving spheres in M are also isotopic to Haken spheres. We now make the following

Claim 5.1. *Suppose that R is a surviving Haken sphere in M , and suppose that R' is a surviving sphere disjoint from R , and not isotopic to R . Then R' is isotopic to a Haken sphere.*

Proof of Claim. Denote by $M - R$ the punctured 3-manifold obtained by cutting at R . $M - R$ has two punctures more than M , corresponding to the two sides of R . $M - R$ has one or two components, depending on whether R is separating or not.

Let M' be the component of $M - R$ containing R' . This manifold inherits a Heegaard splitting from $V \cup_S V'$ with splitting surface a component of $S' = S - (R \cap S)$. If $R \cap S$ is nonseparating, then $g(S') < g(S)$. If $R \cap S$ is separating, then either the genus or the number of boundary components is smaller for S' . In either case, $(g(S'), n(S')) < (g(S), n(S))$ lexicographically.

The sphere R' defines an essential sphere in M' : if it would bound a ball in M' , the same would be true in M (violating incompressibility of R' in M), and if it were isotopic to a boundary component of M' , then R' would be peripheral in M or isotopic to R (both of which we exclude).

Now, if R' is almost peripheral in M' , then by Theorem 3.6 it is isotopic to a Haken sphere in M' . Otherwise, since the complexity of the splitting of M' is smaller than the original one, we can use the inductive hypothesis on M' to conclude that R' is isotopic to a Haken sphere in M' . Interpreting M' as a submanifold of M , and using that the Heegaard

splitting of M' is inherited from M , this shows that R' is isotopic to a Haken sphere in M as well. //

Now, if M contains any surviving spheres, then the Surviving Haken Lemma (Theorem 3.6) implies that there is a surviving Haken sphere σ_0 . Connectivity of the surviving sphere complex (Lemma 3.5), together with the Claim then inductively implies that any surviving sphere is isotopic to a Haken sphere. This finishes the proof of the inductive steps. \square

References

- [1] Bowden, Jonathan and Hensel, Sebastian and Webb, Richard, *Quasi-morphisms on surface diffeomorphism groups*, *J. Amer. Math. Soc.* 35 (2022), no. 1, 211–231.
- [2] Brendle, Tara and Broaddus, Nathan and Putman, Andrew, *The mapping class group of connected sums of $S^2 \times S^1$ s*, preprint
- [3] Griffiths, H. B., **Automorphisms of a 3-dimensional handlebody**. *Abh. Math. Sem. Univ. Hamburg*, 26:191–210, 1963/1964.
- [4] Gültepe, Funda and Leininger, Christoph J and Pho-On, Witsarut, *A universal Cannon-Thurston map and the surviving curve complex*, *Trans. Amer. Math. Soc. Ser. B* 9 (2022), 99–143.
- [5] Haken, Wolfgang, *Some results on surfaces in 3-manifolds*, 1968 Studies in Modern Topology, pp. 39–98, *Math. Assoc. Amer. (distributed by Prentice-Hall, Englewood Cliffs, N.J.)*
- [6] Hatcher, Allen, *Homological stability for automorphism groups of free groups*, *Comment. Math. Helv.* 70 (1995) 39–62.
- [7] Hempel, John, *3-manifolds*, Reprint of the 1976 original. *AMS Chelsea Publishing, Providence, RI*, 2004. xii+195 pp.
- [8] Jaco, William, *Lectures on three-manifold topology*, CBMS Regional Conference Series in Mathematics 43 (1980), *American Mathematical Society, Providence, R.I.*, xii+251 pp.
- [9] Laudenbach, François, *Sur les 2-Sphères D'une Variété de Dimension 3*, *Ann. Math, Second Series*, Vol. 97, No. 1 (1973), 57–81.
- [10] Laudenbach, François, *Topologie de la dimension trois: homotopie et isotopie. (French)*, With an English summary and table of contents. *Astérisque*, No. 12. *Société Mathématique de France, Paris*, 1974. i+152 pp.
- [11] Luft, E., *Actions of the homeotopy group of an orientable 3-dimensional handlebody*. *Math. Ann.*, 234(3):279–292, 1978.
- [12] Oertel, Ulrich and Navarro Carvalho, Leonardo, *A Classification of Automorphisms of Compact 3-Manifolds*, arXiv:math/0510610v1, 2005.

- [13] Qiu, Ruifeng and Scharlemann, Martin, *A proof of the Gordon conjecture*, *Adv. Math.* 222 (2009), no 6, 2085–2106.
- [14] Rolfsen, Dale, *Knots and Links*, Mathematics Lecture Series, No. 7., *Publish or Perish, Inc., Berkeley, Calif.*, 1976. ix+439 pp.
- [15] Scharlemann, Martin, *Generating the genus $g+1$ Goeritz group of a genus g handlebody*, *Geometry and Topology down under*, 347–369, *Contemp. Math.*, 597, *Amer. Math. Soc.*, Providence, RI, 2013.
- [16] Scharlemann, Martin, *A Strong Haken's Theorem*, preprint, arXiv:2003.08523
- [17] Scharlemann, Martin, *Generating the Goeritz Group*, preprint, arXiv:2011.10613
- [18] Schultens, Jennifer *Introduction to 3-manifolds*, Graduate Studies in Mathematics, 151. *American Mathematical Society*, Providence, RI, 2014. x+286 pp.
- [19] Waldhausen, Friedhelm, *Heegaardzerlegungen der 3-Sphäre*, *Topology* 7 (1968), no. 2, 195–203.