

Section 4.1

$$\begin{aligned} 1.) \quad f(x) = 5 &\rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{5 - 5}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = \lim_{h \rightarrow 0} 0 = 0, \\ &\text{so } f'(1) = 0 \end{aligned}$$

$$\begin{aligned} 2.) \quad f(x) = -3x &\rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-3(x+h) - (-3x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{-3x - 3h + 3x}{h} = \lim_{h \rightarrow 0} \frac{-3\cancel{h}}{\cancel{h}} \\ &= \lim_{h \rightarrow 0} -3 = -3 \quad \text{so } f'(-2) = -3 \end{aligned}$$

$$\begin{aligned} 5.) \quad f(x) = 2x^2 &\rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{2(x+h)^2 - 2x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{2(x^2 + 2hx + h^2) - 2x^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{\cancel{2x^2} + 4hx + 2h^2 - \cancel{2x^2}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\cancel{h}(4x + 2h)}{\cancel{h}} = 4x + 2(0) \rightarrow \\ &f'(x) = 4x \quad \text{so } f'(0) = 4(0) = 0 \end{aligned}$$

$$\begin{aligned}
 10.) \quad f(x) &= -x^2 + 4 \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-(x+h)^2 + 4 - (-x^2 + 4)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-(x^2 + 2hx + h^2) + 4 + x^2 - 4}{h} \\
 &= \lim_{h \rightarrow 0} \frac{-x^2 - 2hx - h^2 + x^2}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cancel{h}(-2x - h)}{\cancel{h}} = -2x \text{ so}
 \end{aligned}$$

$$f'(x) = -2x ; \quad f'(x) = 0 \rightarrow -2x = 0 \rightarrow x = 0$$


$$\begin{aligned}
 13.) \quad f(x) &= x^2 - 6x + 9 \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{(x+h)^2 - 6(x+h) + 9 - (x^2 - 6x + 9)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cancel{x^2} + 2hx + h^2 - \cancel{6x} - 6h + 9 - \cancel{x^2} + \cancel{6x} - 9}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cancel{h}(2x + h - 6)}{\cancel{h}} = 2x - 6 \text{ so}
 \end{aligned}$$

$$\begin{aligned}
 f'(x) &= 2x - 6 ; \quad f'(x) = 0 \rightarrow 2x - 6 = 0 \rightarrow \\
 2x &= 6 \rightarrow x = 3 .
 \end{aligned}$$

Section 4.1

21.) a.) $f(x) = 5x^2 \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{5(x+h)^2 - 5x^2}{h} = \lim_{h \rightarrow 0} \frac{5(x^2 + 2hx + h^2) - 5x^2}{h}$
 $= \lim_{h \rightarrow 0} \frac{\cancel{5x^2} + 10hx + 5h^2 - \cancel{5x^2}}{h}$
 $= \lim_{h \rightarrow 0} \frac{\cancel{h}(10x + 5h)}{\cancel{h}} = 10x, \text{ i.e.,}$
 $f'(x) = 10x \rightarrow f'(-1) = 10(-1) = -10$

b.) $f(-1) = 5(-1)^2 = 5(1) = 5$ so $(-1, 5)$ is on the graph; slope of tangent line is $m = f'(-1) = -10$, so line is $y - 5 = -10(x - (-1)) \rightarrow$
 $y - 5 = -10x - 10 \rightarrow y = -10x - 5$



23.) a.) $f(x) = 1 - x^3 \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$
 $= \lim_{h \rightarrow 0} \frac{(1 - (x+h)^3) - (1 - x^3)}{h}$
 $= \lim_{h \rightarrow 0} \frac{1 - (x^3 + 3x^2h + 3xh^2 + h^3) - 1 + x^3}{h}$

$$\begin{aligned}
 &= \lim_{h \rightarrow 0} \frac{\cancel{x-x^3} - 3x^2h - 3xh^2 - h^3 - \cancel{x+x^3}}{h} \\
 &= \lim_{h \rightarrow 0} \frac{h(-3x^2 - 3xh - h^2)}{h} = -3x^2, \text{ i.e.;} \\
 f'(x) &= -3x^2 \rightarrow f'(2) = -3(2)^2 = -12
 \end{aligned}$$

$$\begin{aligned}
 24.) \text{ a.) } f(x) &= \frac{1}{x} \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{\frac{h}{1}} = \lim_{h \rightarrow 0} \frac{x - (x+h)}{(x+h)x} \cdot \frac{1}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\cancel{x-x} - h}{(x+h)x \cdot h} = \lim_{h \rightarrow 0} \frac{-h}{(x+h)x \cdot h} = \frac{-1}{(x)x}, \\
 \text{i.e., } f'(x) &= \frac{-1}{x^2} \rightarrow f'(2) = \frac{-1}{4}
 \end{aligned}$$

$$\begin{aligned}
 25.) f(x) &= \sqrt{x} \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\
 &= \lim_{h \rightarrow 0} \frac{\cancel{(x+h)} - \cancel{x}}{h(\sqrt{x+h} + \sqrt{x})} = \lim_{h \rightarrow 0} \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\
 &= \frac{1}{\sqrt{x} + \sqrt{x}} = \frac{1}{2\sqrt{x}}, \text{ i.e., } f'(x) = \frac{1}{2\sqrt{x}}
 \end{aligned}$$

$$\begin{aligned}
 26.) f(x) &= \frac{1}{x+1} \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\frac{1}{(x+h)+1} - \frac{1}{x+1}}{\frac{h}{1}} = \lim_{h \rightarrow 0} \frac{x+1 - (x+h+1)}{(x+h+1)(x+1)} \cdot \frac{1}{h}
 \end{aligned}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{x+h} - \cancel{x} - h - \cancel{h}}{(x+h+1)(x+1) \cdot h} = \lim_{h \rightarrow 0} \frac{-h}{(x+h+1)(x+1)h}$$

$$= \frac{-1}{(x+1)(x+1)} = \frac{-1}{(x+1)^2}, \text{ i.e., } f'(x) = \frac{-1}{(x+1)^2}$$

30.) $f(x) = x^2 - 3x + 1 \rightarrow f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$$= \lim_{h \rightarrow 0} \frac{[(x+h)^2 - 3(x+h) + 1] - [x^2 - 3x + 1]}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{x^2} + 2hx + h^2 - \cancel{3x} - 3h + 1 - \cancel{x^2} + \cancel{3x} - 1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(2x + h - 3)}{h} = 2x - 3, \text{ i.e.,}$$

$f'(x) = 2x - 3$; pt. $(2, -1)$ so slope of tangent line is $f'(2) = 2(2) - 3 = 1$ and line is $Y - (-1) = (1)(X - 2) \rightarrow Y + 1 = X - 2 \rightarrow Y = X - 3$


36.) If $f(x) = 4x^3$, then

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \rightarrow 0} \frac{4(a+h)^3 - 4a^3}{h}$$

38.) If $f(x) = \sin x$ then

$$f'\left(\frac{\pi}{6}\right) = \lim_{h \rightarrow 0} \frac{f\left(\frac{\pi}{6} + h\right) - f\left(\frac{\pi}{6}\right)}{h} = \lim_{h \rightarrow 0} \frac{\sin\left(\frac{\pi}{6} + h\right) - \sin\frac{\pi}{6}}{h}$$

39.) a.)



$$b.) \text{ ARC} = \frac{s(2) - s(0)}{2 - 0} = \frac{80 - 0}{2} = 40 \text{ km./hr.}$$

$$c.) s'(t) = \lim_{h \rightarrow 0} \frac{s(t+h) - s(t)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{20(t+h)^2 - 20t^2}{h}$$


$$= \lim_{h \rightarrow 0} \frac{20(t^2 + 2th + h^2) - 20t^2}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\cancel{20t^2} + 40th + 20h^2 - \cancel{20t^2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{h(40t + 20h)}{h} = 40t, \text{ i.e.,}$$

$$s'(t) = 40t \quad ; \quad \text{then}$$

$$s'(1) = 40 \text{ km./hr.}$$



b.) i.) $t=0 \rightarrow s=0$

ii.) $t=3 \rightarrow s=0$

iii.) $s'(t) = 3 - 2t = 0 \rightarrow 3 = 2t \rightarrow$

$t = \frac{3}{2}$ sec. (object changes direction) \rightarrow

$s\left(\frac{3}{2}\right) = 3\left(\frac{3}{2}\right) - \left(\frac{3}{2}\right)^2 = \frac{9}{2} - \frac{9}{4} = \frac{9}{4}$ so

particle goes $\frac{9}{4}$ m. right of $s=0$.

iv.) The particle goes infinitely far to the left.

v.) $s'(t) > 0$ for $0 < t < \frac{3}{2}$ sec. ;

$s'(t) = 0$ for $t = \frac{3}{2}$ sec. ;

$s'(t) < 0$ for $t > \frac{3}{2}$ sec.

vi.) velocity of particle is

$s'(t) = 3 - 2t$ m./sec.

49.) $\frac{dN}{dt} = rN\left(1 - \frac{N}{K}\right) = 0 \rightarrow$

$N=0$ (no!) or $1 - \frac{N}{K} = 0 \rightarrow N=K$;

$\frac{dN}{dt} = 0$ means there is zero


growth so maximum capacity has been reached.

51.) a.) False

b.) True


58.)

$$y = 2 - |x - 3|$$




64.) $y = |x^2 - 3|$

not
differentiable
at $x = \sqrt{3}$,
 $x = -\sqrt{3}$




67.) $f(x) = \begin{cases} 2x & , \text{ for } x \leq 1 \\ x+2 & , \text{ for } x > 1 \end{cases}$

not
differentiable
at $x=1$



68.)
$$f(x) = \begin{cases} x^2, & \text{for } x \leq -1 \\ 2 - x^2, & \text{for } x > -1 \end{cases}$$



not differentiable at $x = -1$

Math 17A
 Kouba
 Worksheet 2

Sketch f' from the graph of f .

