


Mean Value Theorem for Integrals : If f is a continuous function on the closed interval $[a, b]$, then there is at least one number c , $a \leq c \leq b$, so that

$$f(c)(b - a) = \int_a^b f(x) dx .$$




Proof : Since f is a continuous function on the closed interval $[a, b]$, by the Maximum- and Minimum-Value Theorems, f has a maximum value M and a minimum value m on $[a, b]$, i.e., $m \leq f(x) \leq M$ on $[a, b]$. By property h of definite integrals,

$$m(b - a) \leq \int_a^b f(x) dx \leq M(b - a) ,$$

so that

$$m \leq \underbrace{\frac{1}{b-a} \int_a^b f(x) dx}_{\text{call this number } y_0} \leq M ,$$




By the Intermediate Value Theorem that

$$f(c) = y_0, \text{ i.e., } f(c) = \frac{1}{b-a} \int_a^b f(x) dx ,$$

so that

$$f(c)(b - a) = \int_a^b f(x) dx .$$


there is at least one number c , $a \leq c \leq b$, so



First Fundamental Theorem of Calculus (FTC1) : Assume that f is a continuous function on the closed interval $[a, b]$ and that $F(x) = \int_a^x f(t) dt$. Then $F'(x) = f(x)$.


Proof : Consider $F(x) = \int_a^x f(t) dt$ as the area under the graph of f above the interval $[a, x]$. Then $F(x+h)$ is the area under the graph of f above the interval $[a, x+h]$ and $F(x+h) - F(x)$ is the area of the "thin strip" from x to $x+h$, i.e., $F(x+h) - F(x) = \int_x^{x+h} f(t) dt$. By the Mean Value Theorem for integrals there is at least one number c , $x \leq c \leq x+h$, so that

$$f(c) \cdot h = \int_x^{x+h} f(t) dt$$



The derivative of $F(x)$ can now be computed as

$$\begin{aligned}
 F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{\int_x^{x+h} f(t) dt}{h} \\
 &= \lim_{h \rightarrow 0} \frac{f(c) h}{h} \\
 &= \lim_{h \rightarrow 0} f(c) \quad (\text{Recall that } x \leq c \leq x+h.) \\
 &= f(x) .
 \end{aligned}$$



Second Fundamental Theorem of Calculus (FTC2) : Let f be a continuous function on the closed interval $[a, b]$. Assume that $F(x)$ is an antiderivative of $f(x)$, i.e., assume that $F'(x) = f(x)$. Then

$$\int_a^b f(x) dx = F(x) \Big|_a^b = F(b) - F(a) .$$

Proof : Let $A(x) = \int_a^x f(t) dt$. Then $A(a) = 0$, $A(b) = \int_a^b f(t) dt$, and $A'(x) = f(x)$ by FTC1. But $F'(x) = f(x)$. By Corollary to the Mean Value Theorem $F(x) = A(x) + C$ for any constant C , or

$$A(x) = F(x) - C .$$

Then

$$\begin{aligned}
 \int_a^b f(x) dx &= \int_a^b f(t) dt \\
 &= A(b) \\
 &= A(b) - A(a) \\
 &= (F(b) - C) - (F(a) - C) \\
 &= F(b) - F(a) \\
 &= F(x) \Big|_a^b .
 \end{aligned}$$