

Section 11.4


$$1.) \sum_{n=1}^{\infty} \frac{x^n}{n^2} ; \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{(n+1)^2} \cdot \frac{n^2}{|x|^n}$$

$$= \lim_{n \rightarrow \infty} |x| \cdot \left(\frac{n}{n+1}\right)^2 = |x| (1)^2 = |x| < 1 \Rightarrow -1 < x < 1;$$

if $x=1$: $\sum_{n=1}^{\infty} \frac{1^n}{n^2} = \sum_{n=1}^{\infty} \frac{1}{n^2}$ is a convergent p -series ($p=2>1$);

if $x=-1$: $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^2}$ converges by the

alternating series test since $a_n = \frac{1}{n^2}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0$; the interval of convergence is $\boxed{-1 \leq x \leq 1}$:




$$2.) \sum_{n=1}^{\infty} \frac{x^n}{\sqrt{n}} ; \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{\sqrt{n+1}} \cdot \frac{\sqrt{n}}{|x|^n}$$

$$= \lim_{n \rightarrow \infty} |x| \sqrt{\frac{n}{n+1}} = |x| \sqrt{1} = |x| < 1 \Rightarrow -1 < x < 1;$$

if $x=1$: $\sum_{n=1}^{\infty} \frac{1^n}{\sqrt{n}} = \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$ diverges by p -series test ($p = \frac{1}{2} \leq 1$); if $x=-1$: $\lim_{n \rightarrow \infty} \sum_{n=1}^{\infty} \frac{(-1)^n}{\sqrt{n}}$

converges by the alternating series test since $a_n = \frac{1}{\sqrt{n}}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}} = 0$; the interval of convergence

is $\boxed{-1 \leq x < 1}$:




3.) $\sum_{n=0}^{\infty} \frac{x^n}{3^n}$; $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{3^{n+1}} \cdot \frac{3^n}{|x|^n}$
 $= \lim_{n \rightarrow \infty} \frac{|x|}{3} = \frac{|x|}{3} < 1 \Rightarrow |x| < 3 \Rightarrow -3 < x < 3$;
 if $x = 3$: $\sum_{n=0}^{\infty} \frac{3^n}{3^n} = \sum_{n=0}^{\infty} 1$ diverges by the
 n th term test; if $x = -3$: $\sum_{n=0}^{\infty} \frac{(-3)^n}{3^n}$
 $= \sum_{n=0}^{\infty} \left(\frac{-3}{3}\right)^n = \sum_{n=0}^{\infty} (-1)^n$ diverges by the
 n th term test since $\lim_{n \rightarrow \infty} (-1)^n$ DNE ($\neq 0$);
 the interval of convergence is $\boxed{-3 < x < 3}$:

4.) $\sum_{n=1}^{\infty} \frac{n^2}{e^n} x^n$; $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{(n+1)^2}{e^{n+1}} |x|^{n+1} \cdot \frac{e^n}{n^2 |x|^n}$
 $= \lim_{n \rightarrow \infty} \left(\frac{n+1}{n}\right)^2 \cdot \frac{|x|}{e} = (1)^2 \frac{|x|}{e} = \frac{|x|}{e} < 1 \Rightarrow |x| < e \Rightarrow$
 $-e < x < e$; if $x = e$: $\sum_{n=1}^{\infty} \frac{n^2}{e^n} \cdot e^n = \sum_{n=1}^{\infty} n^2$
 diverges by the n th term test since
 $\lim_{n \rightarrow \infty} n^2 = \infty \neq 0$; if $x = -e$: $\sum_{n=1}^{\infty} \frac{n^2}{e^n} (-e)^n$
 $= \sum_{n=1}^{\infty} n^2 (-1)^n$ diverges by the n th term
 test since $\lim_{n \rightarrow \infty} n^2 (-1)^n$ DNE ($\neq 0$); the
 interval of convergence is $\boxed{-e < x < e}$:

$$\begin{aligned}
 7.) \quad \sum_{n=0}^{\infty} \frac{x^n}{(2n)!} ; \quad \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} &= \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{(2(n+1))!} \cdot \frac{(2n)!}{|x|^n} \\
 &= \lim_{n \rightarrow \infty} |x| \cdot \frac{\cancel{(2n)}(\cancel{2n-1})(\cancel{2n-2}) \cdots \cancel{3} \cdot \cancel{2} \cdot \cancel{1}}{(2n+2)(2n+1)(\cancel{2n})(\cancel{2n-1}) \cdots \cancel{3} \cdot \cancel{2} \cdot \cancel{1}} \\
 &= \lim_{n \rightarrow \infty} \frac{|x|}{(2n+2)(2n+1)} = 0 < 1 \quad \text{for all values of } x.
 \end{aligned}$$

$$\begin{aligned}
 10.) \quad \sum_{n=0}^{\infty} n! x^n ; \quad \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} &= \lim_{n \rightarrow \infty} \frac{(n+1)! |x|^{n+1}}{n! |x|^n} \\
 &= \lim_{n \rightarrow \infty} (n+1) |x| = \begin{cases} 0 & \text{if } x=0 \\ \infty & \text{if } x \neq 0 \end{cases}, \text{ so series}
 \end{aligned}$$


converges only for $x=0$;





$$\begin{aligned}
 11.) \quad \sum_{n=1}^{\infty} \frac{(-1)^{n+1} x^n}{n} ; \quad \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} &= \lim_{n \rightarrow \infty} \frac{|x|^{n+1}}{n+1} \cdot \frac{n}{|x|^n} \\
 &= \lim_{n \rightarrow \infty} \frac{n}{n+1} \cdot |x| = (1) |x| = |x| < 1 \Rightarrow -1 < x < 1 ;
 \end{aligned}$$

if $x=1$: $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} (1)^n = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$ converges by the alternating series test since $a_n = \frac{1}{n}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$; if $x=-1$: $\sum_{n=1}^{\infty} \frac{(-1)^{n+1} (-1)^n}{n}$
 $= \sum_{n=1}^{\infty} \frac{(-1)^{2n+1}}{n} = \sum_{n=1}^{\infty} \frac{(-1)^{2n} (-1)}{n} = \sum_{n=1}^{\infty} \frac{-1}{n}$ diverges by


the harmonic series test; the interval of convergence is $-1 < x \leq 1$:




12.) $\sum_{n=1}^{\infty} \frac{2^n x^n}{n}$; $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{2^{n+1} |x|^{n+1}}{n+1} \cdot \frac{n}{2^n |x|^n}$
 $= \lim_{n \rightarrow \infty} 2|x| \cdot \frac{n}{n+1} = 2|x|(1) = 2|x| < 1 \Rightarrow |x| < \frac{1}{2} \Rightarrow$
 $-\frac{1}{2} < x < \frac{1}{2}$; if $x = \frac{1}{2}$: $\sum_{n=1}^{\infty} \frac{2^n (\frac{1}{2})^n}{n} = \sum_{n=1}^{\infty} \frac{(2 \cdot \frac{1}{2})^n}{n}$
 $= \sum_{n=1}^{\infty} \frac{1^n}{n} = \sum_{n=1}^{\infty} \frac{1}{n}$ diverges (harmonic series);
 if $x = -\frac{1}{2}$: $\sum_{n=1}^{\infty} \frac{2^n (-\frac{1}{2})^n}{n} = \sum_{n=1}^{\infty} \frac{(2 \cdot -\frac{1}{2})^n}{n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$
 converges by the alternating series test
 since $a_n = \frac{1}{n}$ is \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$; the
 interval of convergence is $\boxed{-\frac{1}{2} \leq x < \frac{1}{2}}$:


14.) $\sum_{n=0}^{\infty} a_n x^n$ (centered at $x=0$), converges at $x=-5$,
 diverges at $x=8$:


- a.) $x=4$: convergence
- b.) $x=4$: absolute convergence
- c.) $x=7$: unknown
- d.) $x=-5$: absolute convergence unknown
- e.) $x=-9$: diverges
- f.) $x=-8$: unknown

15.) $\sum_{n=0}^{\infty} a_n x^n$ (centered at $x=0$), converges for all $x > 0$:
 converges for all $x < 0$.


16.) $\sum_{n=0}^{\infty} a_n 6^n$ converges ($\sum_{n=0}^{\infty} a_n X^n$ centered at $X=0$.):



a.) $X = -6$: unknown

b.) $X = 5$: absolute convergence

c.) $X = -5$: absolute convergence

$$18.) \sum_{n=1}^{\infty} \frac{(x-1)^n}{n 3^n}; \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|x-1|^{n+1}}{(n+1)3^{n+1}} \cdot \frac{n 3^n}{|x-1|^n}$$


$$= \lim_{n \rightarrow \infty} \frac{|x-1|}{3} \cdot \frac{n}{n+1} = \frac{|x-1|}{3} (1) = \frac{|x-1|}{3} < 1 \Rightarrow |x-1| < 3$$

$$\Rightarrow -3 < x-1 < 3 \Rightarrow -2 < x < 4; \text{ if } x=4:$$

$$\sum_{n=1}^{\infty} \frac{3^n}{n 3^n} = \sum_{n=1}^{\infty} \frac{1}{n} \text{ diverges (harmonic series);}$$

$$\text{if } x=-2: \sum_{n=1}^{\infty} \frac{(-3)^n}{n 3^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{ converges by the}$$

alternating series test since $a_n = \frac{1}{n}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$; the interval of convergence is $\boxed{-2 \leq x < 4}$:



$$19.) \sum_{n=0}^{\infty} \frac{(x-1)^n}{n+3}; \lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = \lim_{n \rightarrow \infty} \frac{|x-1|^{n+1}}{n+4} \cdot \frac{n+3}{|x-1|^n}$$


$$= \lim_{n \rightarrow \infty} |x-1| \cdot \frac{n+3}{n+4} = |x-1| (1) = |x-1| < 1 \Rightarrow -1 < x-1 < 1$$

$$\Rightarrow 0 < x < 2; \text{ if } x=0: \sum_{n=0}^{\infty} \frac{(-1)^n}{n+3} \text{ converges by the alternating series test since } a_n = \frac{1}{n+3}$$

is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n+3} = 0$; if $x=2$:

$$\sum_{n=0}^{\infty} \frac{1^n}{n+3} = \sum_{n=0}^{\infty} \frac{1}{n+3} \text{ diverges (harmonic series);}$$

the interval of convergence is $\boxed{0 \leq x < 2}$:



$$22.) \sum_{n=2}^{\infty} \frac{(x-5)^n}{n \ln n} ; \lim_{n \rightarrow \infty} \frac{|x-5|^{n+1}}{(n+1) \ln(n+1)} \cdot \frac{n \ln n}{|x-5|^n}$$

$$= \lim_{n \rightarrow \infty} |x-5| \cdot \frac{n}{n+1} \cdot \frac{\ln n}{\ln(n+1)} = \lim_{n \rightarrow \infty} |x-5| \cdot (1) \cdot \frac{\frac{1}{n}}{\frac{1}{n+1}}$$

$$= \lim_{n \rightarrow \infty} |x-5| \cdot \frac{n+1}{n} = |x-5| (1) = |x-5| < 1 \Rightarrow$$

$$-1 < x-5 < 1 \Rightarrow 4 < x < 6 ; \text{ if } x=4: \sum_{n=2}^{\infty} \frac{(-1)^n}{n \ln n}$$

converges by the alternating series test since $a_n = \frac{1}{n \ln n}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n \ln n} = 0$;


if $x=6$: $\sum_{n=2}^{\infty} \frac{1}{n \ln n} \rightarrow f(x) = \frac{1}{x \ln x}$ is $+$, \downarrow , and

continuous and $\int_2^{\infty} \frac{1}{x \ln x} dx = \lim_{A \rightarrow \infty} \ln |\ln x| \Big|_2^A$

$$= \lim_{A \rightarrow \infty} (\ln |\ln A| - \ln |\ln 2|) = \infty, \text{ so } \sum_{n=2}^{\infty} \frac{1}{n \ln n}$$

diverges by the integral test ; the interval

of convergence is $\boxed{4 \leq x < 6}$;



$$24.) \sum_{n=1}^{\infty} n(x+1)^n ; \lim_{n \rightarrow \infty} \frac{(n+1)|x+1|^{n+1}}{n|x+1|^n} = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right) |x+1|$$

$$= (1) |x+1| < 1 \Rightarrow -1 < x+1 < 1 \Rightarrow -2 < x < 0 ;$$

if $x=0$: $\sum_{n=1}^{\infty} n(1)^n = \sum_{n=1}^{\infty} n$ diverges by the

n th term test since $\lim_{n \rightarrow \infty} n = \infty \neq 0$;

if $x = -2$: $\sum_{n=1}^{\infty} n(-1)^n$ diverges by the n th term test since $\lim_{n \rightarrow \infty} n(-1)^n$ DNE ($\neq 0$); the interval of convergence is

$-2 < x < 0$: $\text{div.} \quad \text{conver.} \quad \text{div.}$
 $x = -2 \quad x = -1 \quad x = 0$
 center

26.) $\sum_{n=0}^{\infty} (-1)^n \frac{(x+4)^n}{n+2}$; $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|}$

$= \lim_{n \rightarrow \infty} \frac{|x+4|^{n+1}}{n+3} \cdot \frac{n+2}{|x+4|^n} = \lim_{n \rightarrow \infty} \frac{1+\frac{2}{n}}{1+\frac{3}{n}} |x+4|$

$= (1) |x+4| = |x+4| < 1 \Rightarrow -1 < x+4 < 1 \Rightarrow -5 < x < -3$;

if $x = -3$: $\sum_{n=0}^{\infty} (-1)^n \frac{1^n}{n+2} = \sum_{n=0}^{\infty} (-1)^n \cdot \frac{1}{n+2}$ converges

by the alternating series test since $a_n = \frac{1}{n+2}$ is $+$, \downarrow , and $\lim_{n \rightarrow \infty} \frac{1}{n+2} = 0$; if $x = -5$:

$\sum_{n=0}^{\infty} (-1)^n \frac{(-1)^n}{n+2} = \sum_{n=0}^{\infty} (-1)^{2n} \cdot \frac{1}{n+2} = \sum_{n=0}^{\infty} ((-1)^2)^n \cdot \frac{1}{n+2}$
 $= \sum_{n=0}^{\infty} 1^n \cdot \frac{1}{n+2} = \sum_{n=0}^{\infty} \frac{1}{n+2}$ diverges by the

harmonic series test; the interval of convergence is $-5 < x \leq -3$:


$\text{div.} \quad \text{conver.} \quad \text{div.}$
 $x = -5 \quad x = -4 \quad x = -3$
 center

33.) a.) $\sum_{n=0}^{\infty} a_n x^5$ (centered at $x = 0$) diverges

at $x = 3$: $\text{div.} \quad \text{div.}$
 $x = -3 \quad x = 0 \quad x = 3$
 center


series must diverge at $x \geq 3$ and $x < -3$.

b.) $\sum_{n=0}^{\infty} a_n (x+5)^n$ (centered at $x = -5$) diverges



series must diverge for $x \geq -2$ and $x < -8$.

34.) $\sum_{n=0}^{\infty} a_n (x-3)^n$ (centered at $x = 3$) converges



series must converge for $-1 < x \leq 7$.