## Math 21D

## Vogler

## Discussion Sheet 5

- 1.) Show that the curve plotted by projectile motion,  $\vec{r}(t) = (|\vec{v}_0| \cos \alpha \cdot t)\vec{i} + (|\vec{v}_0| \sin \alpha \cdot t (1/2)gt^2)\vec{j}$ , is a parabola in the *xy*-plane.
- 2.) Show that the maximum downrange distance for projectile motion with a given initial speed  $|\vec{v}_0|$  occurs when  $\alpha=45^o$  and is  $x=\frac{|\vec{v}_0|^2}{g}$  (Hint: See formula III on Projectile Motion Handout.).
- 3.) Consider path C plotted by vector function  $\vec{r}(t) = t \vec{i} + \sqrt{t} \vec{j}$  for  $0 \le t \le 9$ .
  - a.) Sketch C.
  - b.) Find  $\vec{v}(t)$ ,  $\vec{T}(t)$ ,  $\vec{N}(t)$ , and  $\vec{a}(t)$ .
- c.) Plot  $\vec{r}(1), \vec{v}(1), \vec{T}(1), \vec{N}(1),$  and  $\vec{a}(1)$ . Also compute the speed and acceleration of motion when t=1.
- 4.) Let  $\vec{r}(t) = (12\sin t) \vec{i} + (-12\cos t) \vec{j} + (5t) \vec{k}$  be a position vector function.
  - a.) Determine the position vector when
    - i.) t = 0 ii.)  $t = 7\pi/6$
- b.) Write t as a function of arc length s. Write  $\vec{r}(t)$  as a function of arc length s, i.e., write  $\vec{r}(t) = \vec{r}(t(s))$ .
  - c.) Determine the position vector when
    - i.) s = 0 ii.)  $s = 39\pi$
- 5.) Determine the length of path C determined  $\vec{r}(t) = (\cos^3 t) \ \vec{i} + (\sin^3 t) \ \vec{j}$  for  $0 \le t \le 2\pi$ .
- 6.) Evaluate the following line integrals.

a.) 
$$\int_C 2x \ ds$$
 ,  $\vec{r}(t) = (1/2)t^2 \ \vec{i} + (1/4)t^4 \ \vec{j}$  for  $0 \le t \le 2$ 

b.) 
$$\int_C \sqrt{x^2 + z^2} \ ds$$
,  $\vec{r}(t) = (2\cos t) \ \vec{j} + (2\sin t) \ \vec{k}$  for  $\pi \le t \le 2\pi$ 

c.) 
$$\int_C 3xyz \ ds$$
 ,  $\vec{r}(t) = t \ \vec{i} + 2t \ \vec{i} - t \ \vec{j}$  for  $0 \le t \le 4$ 

- 7.) A spring lies on the path determined by  $\vec{r}(t) = (\sin t) \ \vec{i} + (\cos t) \ \vec{j} + (2/3)t^{3/2} \ \vec{k}$  for  $0 \le t \le 4\pi$ . Sketch the wire and find it's length.
- 8.) Find the area of the vertical wall sitting on the xy-plane on the line y=2x from x=0 to x=4, if the height of the wall at the point (x,y) is  $xy^2$ .
- 9.) A wire lies on the path determined by the helix  $\vec{r}(t) = \cos t \ \vec{i} + \sin t \ \vec{j} + t \ \vec{k}$  for  $0 \le t \le 2\pi$ . It's density at point (x, y, z) is given by  $\delta(x, y, z) = xy + z + 3$ . Compute the
  - a.) length of the wire.
  - b.) mass of the wire.
  - c.) x-coordinate for its center of mass.
  - d.) z-coordinate for its centroid.
  - e.) moment of the wire relative to the plane y = 1.
  - f.) moment of inertia of the wire about
    - i.) the origin
- ii.) the z-axis

## THE FOLLOWING PROBLEM IS FOR RECREATIONAL PURPOSES ONLY.

10.) Divide the following figure into 4 parts each of the same size and shape.

