

Fall 2003 Mathematics Graduate Program MA Exam

Instructions: Explain your answers clearly. Unclear answers will not receive credit. State results and theorems that you are using.

1. ANALYSIS

Problem 1. (a) For a function $f : (a, b) \rightarrow \mathbb{R}^1$, (a, b) an open interval, state briefly but precisely:

- What is meant by the statement: $f(x)$ is continuous at $x_0 \in (a, b)$.
- What is meant by the statement: $f(x)$ is continuous on (a, b) .
- What is meant by the statement: $f(x)$ is uniformly continuous on (a, b) .

(b) Prove, directly from the definition, that the function $f(x) = 1/x$ is uniformly continuous on the interval $[1, \infty)$.

Problem 2. Let $\{U_n\}_{n=1}^\infty$ be a nested sequence of open sets in a topological space X , so that $U_1 \subset U_2 \subset \dots \subset U_n \subset U_{n+1}$. Let $x_n \in U_n \setminus U_{n-1}$. Set $U = \bigcup_{n=1}^\infty U_n$. Prove that $\{x_n\}$ does not have a subsequence that converges to a point in U .

Problem 3. Let $T : (X, d) \rightarrow (X, d)$ be a contraction mapping from the metric space (X, d) to itself, so that for some $r < 1$, $d(Tx, Ty) \leq rd(x, y) \forall x, y \in X$. Assume that x_0 is a fixed point of this mapping. Prove that

$$d(x, x_0) \leq \frac{d(x, T(x))}{1-r}$$

Problem 4. Let y, y' be two elements of a Hilbert space H . Prove that if $\langle y, x \rangle = \langle y', x \rangle$ for every $x \in H$ then $y = y'$.

Problem 5. Let L and R be the left shift operator and the right shift operator of $l^2(\mathbb{N})$ respectively. So

$$\begin{aligned} L(x_1, x_2, x_3, \dots) &= (x_2, x_3, x_4, \dots) \\ R(x_1, x_2, x_3, \dots) &= (0, x_1, x_2, x_3, x_4, \dots). \end{aligned}$$

Find the point spectrum of L and R

Problem 6. Define the following three sequences of functions $[0, +\infty) \rightarrow \mathbb{R}$:

$$\begin{aligned} (f_n)_{n=1}^\infty \text{ given by } f_n(x) &= \begin{cases} \frac{n^{1/2}}{(x+1)^n} & \text{if } 0 \leq x \leq n \\ 0 & \text{else} \end{cases} \\ (g_n)_{n=1}^\infty \text{ given by } g_n(x) &= \begin{cases} \sin(2\pi nx) & \text{if } n \leq x \leq n+1 \\ 0 & \text{else} \end{cases} \end{aligned}$$

$$\text{and } (h_n)_{n=1}^\infty \text{ given by } h_n(x) = \sum_{k=1}^n \frac{k}{\sqrt{n}} \text{Ind}_{[k, k+(1/n^2)]}(x).$$

Consider these sequences with each of the topologies given below and determine whether or not they converge and, if they converge, determine their limits. Explain your assertions.

- Pointwise on $[0, +\infty)$.
- Uniformly on $[0, +\infty)$.
- In the norm topology of $L^2([0, +\infty))$.
- Strongly in $L^{3/2}([0, +\infty))$.
- Weakly in $L^{3/2}([0, +\infty))$.

Fall 2003 Mathematics Graduate Program MA Exam

2. ALGEBRA AND LINEAR ALGEBRA

Problem 7. Let G be a group and p a prime. Prove or give a counter example:

- a.** A group of order p is commutative.
- b.** A group of order p^2 is commutative.
- c.** A group of order p^3 is commutative.

Problem 8. Let F be a finite field. Show that the number of elements of F is p^r for some prime p and positive integer r .

Problem 9. A vector space V contains an n -element set with the following properties:

- (i) It is not linearly independent, but contains an $(n - 1)$ -element linearly independent set;
- (ii) It does not span V , but is contained in an $(n + 1)$ -element spanning set.

Prove that $\dim V = n$.

Problem 10. Let B be a symmetric, non-degenerate, not positive definite bilinear form in an n -dimensional real vector space V . Prove that there exists a basis v_1, \dots, v_n in V such that $B(v_i, v_i) < 0$ for all i .

Problem 11. Let $I \subset \mathbb{R}[x]$ be the ideal generated by the polynomial $x^2 + 2x + 3$. Prove that the quotient ring $\mathbb{R}[x]/I$ is isomorphic to the field \mathbb{C} of complex numbers.

Problem 12. Let $K = \mathbb{Q}(\sqrt{2}, \sqrt{3})$. Find α so that $K = \mathbb{Q}(\alpha)$, and compute the irreducible polynomial of α over \mathbb{Q} .