

Linearity Principles

Recall the example of the homogeneous linear equation from the end of last class.

Example. $\frac{dy}{dt} = \frac{-ty}{1+t^2}$

We derived the general solution $y(t) = \frac{k}{\sqrt{1+t^2}}$ where k is an arbitrary constant.



Linearity Principles

Why are linear equations so much more amenable to analytic techniques than nonlinear equations? The reason is that their solutions satisfy important linearity principles.

Let's begin with homogeneous linear equations:

Linearity Principle. If $y_h(t)$ is a solution of a homogeneous linear differential equation

$$\frac{dy}{dt} = a(t)y,$$

then any *constant* multiple $y_k(t) = ky_h(t)$ of $y_h(t)$ is also a solution. In other words, given a constant $k \neq 1$ and a solution $y_h(t)$, we obtain another solution by multiplying $y_h(t)$ by k .

Note that the Linearity Principle is not true for nonlinear equations. For example, consider the nonlinear equation

$$\frac{dy}{dt} = y^2.$$

You can check that the function

$$y_1(t) = \frac{1}{1-t}$$

is a solution but that

$$y_2(t) = 2y_1(t) = \frac{2}{1-t}$$

is not a solution.

There is a similar “linearity” principle for nonhomogeneous linear equations:

Extended Linearity Principle For First-Order Equations. Consider a first-order, nonhomogeneous, linear equation

$$\frac{dy}{dt} = a(t)y + b(t)$$

and its associated homogeneous equation

$$\frac{dy}{dt} = a(t)y.$$


1. If $y_h(t)$ is any solution of the homogeneous equation and $y_p(t)$ (“ p ” for particular solution) is *any* solution of the nonhomogeneous equation, then $y_h(t) + y_p(t)$ is also a solution of the nonhomogeneous equation.
2. Suppose $y_p(t)$ and $y_q(t)$ are two solutions of the nonhomogeneous equation. Then $y_p(t) - y_q(t)$ is a solution of the associated homogeneous equation.

Therefore, if $y_h(t)$ is nonzero, $ky_h(t) + y_p(t)$ is the general solution of the nonhomogeneous equation.

We can paraphrase the Extended Linearity Principle by saying that:

The general solution of a nonhomogeneous linear equation consists of the sum of *any* particular solution of the nonhomogeneous equation and the general solution of the associated homogeneous equation.

Example. $\frac{dy}{dt} = \frac{-ty}{1+t^2} + \frac{2t^2+1}{4t^2+4}$



Example 2. $\frac{dy}{dt} = -y + 2 \cos 4t$

1. General solution of the associated homogeneous equation:

2. Particular solution of the nonhomogeneous equation:

