

---

---

More general comments

At the end of last class, I started to make some general comments about first-order differential equations, and I want to continue with those comments now.

1. What does it mean to solve the initial-value problem

$$\frac{dy}{dt} = f(t, y), \quad y(t_0) = y_0?$$

A solution to the initial-value problem is a differentiable function  $y(t)$  defined on some interval  $a < t_0 < b$  containing  $t_0$  such that

(a)  $y(t_0) = y_0$  and

(b)  $\frac{dy}{dt} = f(t, y(t))$  for all  $t$  in the interval  $a < t < b$ .

2. Be careful about notation: The distinction between the independent and the dependent variables is important.

**Example 1.**  $\frac{dy}{dt} = kt$

The solutions to this equation are  $y(t) = k\frac{t^2}{2} + c$ , where  $c$  is an arbitrary constant.

**Example 2.**  $\frac{dy}{dt} = ky$

The solutions to this equation are  $y(t) = y_0e^{kt}$ , where  $y_0$  is an arbitrary constant.

3. What does the term **general solution** mean?

---

---


4. You should never get a wrong answer in this course:

- 
- 
5. Even relatively simple looking differential equations can have solutions that cannot be expressed in terms of functions that we already know and love.

Consider the initial-value problem

$$\frac{dy}{dt} = y^3 + t^2, \quad y(0) = 0.$$

Here is the graph of the solution as generated by HPGSolver.



Our general approach in this course:

We will study differential equations

1. using the theory and
2. various techniques:
  - (a) analytic techniques
  - (b) geometric/qualitative techniques, and
  - (c) numerical techniques.

---

---

## Separable Differential Equations (an analytic technique)

First let's recall the method of substitution for calculating integrals (really antiderivatives):

---

---

A differential equation

$$\frac{dy}{dt} = f(t, y)$$

is **separable** if it can be written in the form

$$\frac{dy}{dt} =$$

**Two Examples:**

1.  $\frac{dy}{dt} = -2ty^2$

2.  $\frac{dy}{dt} = y^3 + t^2$

---

---


Let's go back to the first example

**Example.**  $\frac{dy}{dt} = -2ty^2$

(Additional blank space on top of next page.)



We turn to `FirstOrderExamples` to get a sense of the graphs of these solutions:



What's the general solution to  $\frac{dy}{dt} = -2ty^2$ ? (Think before you answer.)