

### PREREQUISITE REVIEW 5.1

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, rewrite the expression using rational exponents.

1.  $\frac{\sqrt{x}}{x}$

2.  $\sqrt[3]{2x}(2x)$

3.  $\sqrt{5x^3} + \sqrt{x^5}$

4.  $\frac{1}{\sqrt{x}} + \frac{1}{\sqrt[3]{x^2}}$

5.  $\frac{(x+1)^3}{\sqrt{x+1}}$

6.  $\frac{\sqrt{x}}{\sqrt[3]{x}}$

In Exercises 7–10, let  $(x, y) = (2, 2)$ , and solve the equation for  $C$ .

7.  $y = x^2 + 5x + C$

8.  $y = 3x^3 - 6x + C$

9.  $y = -16x^2 + 26x + C$

10.  $y = -\frac{1}{4}x^4 - 2x^2 + C$

### EXERCISES 5.1

In Exercises 1–8, verify the statement by showing that the derivative of the right side is equal to the integrand of the left side.

1.  $\int \left(-\frac{9}{x^4}\right) dx = \frac{3}{x^3} + C$

2.  $\int \frac{4}{\sqrt{x}} dx = 8\sqrt{x} + C$

3.  $\int \left(4x^3 - \frac{1}{x^2}\right) dx = x^4 + \frac{1}{x} + C$

4.  $\int \left(1 - \frac{1}{\sqrt[3]{x^2}}\right) dx = x - 3\sqrt[3]{x} + C$

5.  $\int 2\sqrt{x}(x-3) dx = \frac{4x^{3/2}(x-5)}{5} + C$

6.  $\int 4\sqrt{x}(x^2-2) dx = \frac{8x^{3/2}(3x^2-14)}{21} + C$

7.  $\int \frac{x^2-1}{x^{3/2}} dx = \frac{2(x^2+3)}{3\sqrt{x}} + C$

8.  $\int \frac{2x-1}{x^{4/3}} dx = \frac{3(x+1)}{\sqrt[3]{x}} + C$

In Exercises 9–20, find the indefinite integral and check your result by differentiation.

9.  $\int 6 dx$

10.  $\int -4 dx$

11.  $\int 5t^2 dt$

12.  $\int 3t^4 dt$

13.  $\int 5x^{-3} dx$

14.  $\int 4y^{-3} dy$

15.  $\int du$

16.  $\int dr$

17.  $\int e dt$

18.  $\int e^3 dy$

19.  $\int y^{3/2} dy$

20.  $\int v^{-1/2} dv$

In Exercises 21–28, find the indefinite integral using the column in Example 4 as a model. Use a symbolic integration utility to verify your results.

21.  $\int \sqrt[3]{x} dx$

22.  $\int \frac{1}{x^2} dx$

23.  $\int \frac{1}{x\sqrt{x}} dx$

24.  $\int \frac{1}{x^2\sqrt{x}} dx$

25.  $\int x(x^2+3) dx$


26.  $\int t(t^2+2) dt$

27.  $\int \frac{1}{2x^3} dx$


28.  $\int \frac{1}{8x^3} dx$

In Exercises 29–32, find two functions that have the given derivative, and sketch the graph of each. (There is more than one correct answer.)


29.



30.



31.



In Exercises 33–36, result by differentiation.

33.  $\int (x^3 + 2)$

35.  $\int (\sqrt[3]{x} - \frac{1}{x})$

37.  $\int (\sqrt[3]{x^2} +$

39.  $\int \frac{1}{3x^4} dx$

41.  $\int \frac{2x^3 + 1}{x^3}$

In Exercises 43–46, find the indefinite integral.

43.  $\int u(3u^2 +$

45.  $\int (x-1)($

47.  $\int y^2\sqrt{y} dy$

In Exercises 49–54, find the derivative of the function.

49.  $f(x) = 3\sqrt{x}$

50.  $f(x) = \frac{1}{3}x$

51.  $f(x) = 6x^2$



52.  $f(x) = (2x - 1)^2$

53.  $f(x) = \frac{2}{x}$

54.  $f(x) = \frac{x^2}{x-1}$

In Exercises 55–58, find the general equation of the line tangent to the curve at the given point.

55.  $f(x) = -5x$



In Exercises 33–42, find the indefinite integral and check your result by differentiation.

33.  $\int (x^3 + 2) dx$

34.  $\int (x^2 - 2x + 3) dx$

35.  $\int \left( \sqrt[3]{x} - \frac{1}{2\sqrt[3]{x}} \right) dx$

36.  $\int \left( \sqrt{x} + \frac{1}{2\sqrt{x}} \right) dx$

37.  $\int (\sqrt[3]{x^2} + 1) dx$

38.  $\int (\sqrt[4]{x^3} + 1) dx$

39.  $\int \frac{1}{3x^4} dx$

40.  $\int \frac{1}{4x^2} dx$

41.  $\int \frac{2x^3 + 1}{x^3} dx$

42.  $\int \frac{t^2 + 2}{t^2} dt$

In Exercises 43–48, use a symbolic integration utility to find the indefinite integral.

43.  $\int u(3u^2 + 1) du$

44.  $\int \sqrt{x}(x + 1) dx$

45.  $\int (x - 1)(6x - 5) dx$

46.  $\int (2t^2 - 1)^2 dt$

47.  $\int y^2 \sqrt{y} dy$

48.  $\int (1 + 3t)t^2 dt$

In Exercises 49–54, find the particular solution  $y = f(x)$  that satisfies the differential equation and initial condition.

49.  $f'(x) = 3\sqrt{x} + 3$ ;  $f(1) = 4$

50.  $f'(x) = \frac{1}{5}x - 2$ ;  $f(10) = -10$

51.  $f'(x) = 6x(x - 1)$ ;  $f(1) = -1$

52.  $f'(x) = (2x - 3)(2x + 3)$ ;  $f(3) = 0$

53.  $f'(x) = \frac{2-x}{x^3}$ ,  $x > 0$ ;  $f(2) = \frac{3}{4}$

54.  $f'(x) = \frac{x^2 - 5}{x^2}$ ,  $x > 0$ ;  $f(1) = 2$

In Exercises 55 and 56, you are shown a family of graphs, each of which is a general solution of the given differential equation. Find the equation of the particular solution that passes through the indicated point.

55.  $\frac{dy}{dx} = -5x - 2$

56.  $\frac{dy}{dx} = 2(x - 1)$

Figure for 55





Figure for 56



In Exercises 57 and 58, find the equation of the function  $f$  whose graph passes through the point.

Derivative	Point
57. $f'(x) = 6\sqrt{x} - 10$	(4, 2)

58. $f'(x) = \frac{6}{x^2}$	(2, 5)
-----------------------------	--------

In Exercises 59–62, find a function  $f$  that satisfies the conditions.

59.  $f''(x) = 2$ ,  $f'(2) = 5$ ,  $f(2) = 10$

60.  $f''(x) = x^2$ ,  $f'(0) = 6$ ,  $f(0) = 3$

61.  $f''(x) = x^{-2/3}$ ,  $f'(8) = 6$ ,  $f(0) = 0$

62.  $f''(x) = x^{-3/2}$ ,  $f'(1) = 2$ ,  $f(9) = -4$

**Cost** In Exercises 63–66, find the cost function for the marginal cost and fixed cost.

Marginal Cost	Fixed Cost ( $x = 0$ )
63. $\frac{dC}{dx} = 85$	\$5500

64. $\frac{dC}{dx} = \frac{1}{50}x + 10$	\$1000
--	--------

65. $\frac{dC}{dx} = \frac{1}{20\sqrt{x}} + 4$	\$750
--	-------

66. $\frac{dC}{dx} = \frac{\sqrt[4]{x}}{10} + 10$	\$2300
---	--------

**Demand Function** In Exercises 67–70, find the revenue and demand functions for the given marginal revenue. (Use the fact that  $R = 0$  when  $x = 0$ .)

67.  $\frac{dR}{dx} = 225 - 3x$

68.  $\frac{dR}{dx} = 310 - 4x$

69.  $\frac{dR}{dx} = 225 + 2x - x^2$

70.  $\frac{dR}{dx} = 100 - 6x - 2x^2$

**Profit** In Exercises 71–74, find the profit function for the given marginal profit and initial condition.

Marginal Profit      Initial Condition

71.  $\frac{dP}{dx} = -18x + 1650$        $P(15) = \$22,725$

72.  $\frac{dP}{dx} = -40x + 250$        $P(5) = \$650$

73.  $\frac{dP}{dx} = -24x + 805$        $P(12) = \$8000$

74.  $\frac{dP}{dx} = -30x + 920$        $P(8) = \$6500$

**Vertical Motion** In Exercises 75–78, use  $a(t) = -32$  feet per second per second as the acceleration due to gravity.

75. A ball is thrown vertically upward with an initial velocity of 60 feet per second. How high will the ball go?

76. The Grand Canyon is 6000 feet deep at the deepest part. A rock is dropped from this height. Express the height of the rock as a function of the time  $t$  (in seconds). How long will it take the rock to hit the canyon floor?

77. With what initial velocity must an object be thrown upward from the ground to reach the height of the Washington Monument (550 feet)?

78. A balloon, rising vertically with a velocity of 16 feet per second, releases a sandbag at an instant when the balloon is 64 feet above the ground.

(a) How many seconds after its release will the bag strike the ground?

(b) With what velocity will the bag strike the ground?

79. **Cost** A company produces a product for which the marginal cost of producing  $x$  units is modeled by

$$\frac{dC}{dx} = 2x - 12$$

and the fixed costs are \$125.

(a) Find the total cost function and the average cost function.

(b) Find the total cost of producing 50 units.

(c) In part (b), how much of the total cost is fixed? How much is variable? Give examples of fixed costs associated with the manufacturing of a product. Give examples of variable costs.

80. **Population Growth** The growth rate of Horry County in South Carolina can be modeled by

$$\frac{dP}{dt} = 105.74t + 2639.3$$

where  $t$  is the time in years, with  $t = 0$  corresponding to 1970. The county's population was 196,629 in 2000.

(a) Find the model for Horry County's population.

(b) Use the model to predict the population in 2010. Does your answer seem reasonable? Explain your reasoning.

81. **Vital Statistics** The rate of increase of the number of married couples  $M$  (in thousands) in the United States from 1970 to 2000 can be modeled by

$$\frac{dM}{dt} = 0.636t^2 - 28.48t + 632.7$$

where  $t$  is the time in years, with  $t = 0$  corresponding to 1970. The number of married couples in 2000 was 56,497 thousand. (Source: U.S. Census Bureau)

(a) Find the model for the number of married couples in the United States.

(b) Use the model to predict the number of married couples in the United States in 2010. Does your answer seem reasonable? Explain your reasoning.

82. **Economics: Marginal Benefits and Costs** The table gives the marginal benefit and marginal cost of producing  $x$  products for a given company. Plot the points in each column and use the *regression* feature of a graphing utility to find a linear model for marginal benefit and a quadratic model for marginal cost. Then use integration to find the benefit  $B$  and cost  $C$  equations. Assume  $B(0) = 0$  and  $C(0) = 425$ . Finally, find the intervals in which the benefit exceeds the cost of producing  $x$  products, and make a recommendation for how many products the company should produce based on your findings. (Source: Adapted from *Baylan Economics, Fourth Edition*)

Number of products	1	2	3	4	5
Marginal benefit	330	320	290	270	250
Marginal cost	150	120	100	110	120

Number of products	6	7	8	9	10
Marginal benefit	230	210	190	170	160
Marginal cost	140	160	190	250	320

83. **Research Project** Use your school's library, the Internet, or some other reference source to research a company that markets a natural resource. Find data on the revenue of the company and on the consumption of the resource. Then find a model for each. Is the company's revenue related to the consumption of the resource? Explain your reasoning.

## The Ge

In Section

$$\int x^n dx$$

to find ant  
you will  
functions.

To be  
Because y  
might disc

$$\frac{d}{dx} [C$$

$$\frac{d}{dx} [(x$$

$$(x^2 +$$

The key to  
words, thi  
Letting  $u$

$$\int (x^2$$

This is an

## Gener

If  $u$  is

$$\int$$

When  
integrand  
is also a fi

**PREREQUISITE  
REVIEW 5.2**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–10, find the indefinite integral.

1.  $\int (2x^3 + 1) dx$

2.  $\int (x^{1/2} + 3x - 4) dx$

3.  $\int \frac{1}{x^2} dx$

4.  $\int \frac{1}{3t^3} dt$

5.  $\int (1 + 2t)t^{3/2} dt$

6.  $\int \sqrt{x}(2x - 1) dx$

7.  $\int \frac{5x^3 + 2}{x^2} dx$

8.  $\int \frac{2x^2 - 5}{x^4} dx$

9.  $\int (x^2 + 1)^2 dx$

10.  $\int (x^3 - 2x + 1)^2 dx$

In Exercises 11–14, simplify the expression.

11.  $\left(-\frac{5}{4}\right)\frac{(x-2)^4}{4}$

12.  $\left(\frac{1}{6}\right)\frac{(x-1)^{-2}}{-2}$

13.  $(6)\frac{(x^2+3)^{2/3}}{2/3}$

14.  $\left(\frac{5}{2}\right)\frac{(1-x^3)^{-1/2}}{-1/2}$

**EXERCISES 5.2**

In Exercises 1–8, identify  $u$  and  $du/dx$  for the integral  $\int u^n(du/dx) dx$ .

1.  $\int (5x^2 + 1)^2(10x) dx$

2.  $\int (3 - 4x^2)^3(-8x) dx$

3.  $\int \sqrt{1-x^2}(-2x) dx$

4.  $\int 3x^2\sqrt{x^3+1} dx$

5.  $\int \left(4 + \frac{1}{x^2}\right)^5\left(\frac{-2}{x^3}\right) dx$

6.  $\int \frac{1}{(1+2x)^2}(2) dx$

7.  $\int (1 + \sqrt{x})^3\left(\frac{1}{2\sqrt{x}}\right) dx$

8.  $\int (4 - \sqrt{x})^2\left(\frac{-1}{2\sqrt{x}}\right) dx$

19.  $\int \frac{x+1}{(x^2+2x-3)^2} dx$

20.  $\int \frac{6x}{(1+x^2)^3} dx$

21.  $\int \frac{x-2}{\sqrt{x^2-4x+3}} dx$

22.  $\int \frac{4x+6}{(x^2+3x+7)^3} dx$

23.  $\int 5u\sqrt[3]{1-u^2} du$

24.  $\int u^3\sqrt{u^4+2} du$

25.  $\int \frac{4y}{\sqrt{1+y^2}} dy$

26.  $\int \frac{x^2}{\sqrt{1-x^3}} dx$

27.  $\int \frac{-3}{\sqrt{2t+3}} dt$

28.  $\int \frac{t+2t^2}{\sqrt{t}} dt$

In Exercises 9–28, find the indefinite integral and check the result by differentiation.

9.  $\int (1+2x)^4(2) dx$

10.  $\int (x^2-1)^3(2x) dx$

11.  $\int \sqrt{5x^2-4}(10x) dx$

12.  $\int \sqrt{3-x^3}(3x^2) dx$

13.  $\int (x-1)^4 dx$


14.  $\int (x-3)^{5/2} dx$

15.  $\int x(x^2-1)^7 dx$

16.  $\int x(1-2x^2)^3 dx$

17.  $\int \frac{x^2}{(1+x^3)^2} dx$

18.  $\int \frac{x^2}{(x^3-1)^2} dx$

 In Exercises 29–34, use a symbolic integration utility to find the indefinite integral.

29.  $\int \frac{x^3}{\sqrt{1-x^4}} dx$

30.  $\int \frac{3x}{\sqrt{1-4x^2}} dx$

31.  $\int \left(1 + \frac{4}{t^2}\right)^2\left(\frac{1}{t^3}\right) dt$

32.  $\int \left(1 + \frac{1}{t}\right)^3\left(\frac{1}{t^2}\right) dt$

33.  $\int (x^3+3x)(x^2+1) dx$

34.  $\int (3-2x-4x^2)(1+4x) dx$

In Exercises 35–42, use formal substitution (as illustrated in Example 5) to find the indefinite integral.

35.  $\int x(6x^2 - 1)^3 dx$       36.  $\int x^2(1 - x^3)^2 dx$   
 37.  $\int x^2(2 - 3x^3)^{3/2} dx$       38.  $\int t\sqrt{t^2 + 1} dt$   
 39.  $\int \frac{x}{\sqrt{x^2 + 25}} dx$       40.  $\int \frac{3}{\sqrt{2x + 1}} dx$   
 41.  $\int \frac{x^2 + 1}{\sqrt{x^3 + 3x + 4}} dx$       42.  $\int \sqrt{x}(4 - x^{3/2})^2 dx$

In Exercises 43–46, (a) perform the integration in two ways: once using the Simple Power Rule and once using the General Power Rule. (b) Explain the difference in the results. (c) Which method do you prefer? Explain your reasoning.

43.  $\int (2x - 1)^2 dx$       44.  $\int (3 - 2x)^2 dx$   
 45.  $\int x(x^2 - 1)^2 dx$       46.  $\int x(2x^2 + 1)^2 dx$

47. Find the equation of the function  $f$  whose graph passes through the point  $(0, \frac{4}{3})$  and whose derivative is  $f'(x) = x\sqrt{1 - x^2}$ .

48. Find the equation of the function  $f$  whose graph passes through the point  $(0, \frac{7}{3})$  and whose derivative is  $f'(x) = x\sqrt{1 - x^2}$ .

49. **Cost** The marginal cost of a product is modeled by

$$\frac{dC}{dx} = \frac{4}{\sqrt{x + 1}}$$

When  $x = 15$ ,  $C = 50$ .

- (a) Find the cost function.  
 (b) Use a graphing utility to graph  $dC/dx$  and  $C$  in the same viewing window.

50. **Cost** The marginal cost of a product is modeled by

$$\frac{dC}{dx} = \frac{12}{\sqrt[3]{12x + 1}}$$

When  $x = 13$ ,  $C = 100$ .

- (a) Find the cost function.  
 (b) Use a graphing utility to graph  $dC/dx$  and  $C$  in the same viewing window.

**Supply Function** In Exercises 51 and 52, find the supply function  $x = f(p)$  that satisfies the initial conditions.

51.  $\frac{dx}{dp} = p\sqrt{p^2 - 25}$ ,  $x = 600$  when  $p = \$13$

52.  $\frac{dx}{dp} = \frac{10}{\sqrt{p - 3}}$ ,  $x = 100$  when  $p = \$3$

**Demand Function** In Exercises 53 and 54, find the demand function  $x = f(p)$  that satisfies the initial conditions.

53.  $\frac{dx}{dp} = -\frac{6000p}{(p^2 - 16)^{3/2}}$ ,  $x = 5000$  when  $p = \$5$

54.  $\frac{dx}{dp} = -\frac{400}{(0.02p - 1)^3}$ ,  $x = 10,000$  when  $p = \$100$

55. **Gardening** An evergreen nursery usually sells a type of shrub after 5 years of growth and shaping. The growth rate during those 5 years is approximated by

$$\frac{dh}{dt} = \frac{17.6t}{\sqrt{17.6t^2 + 1}}$$

where  $t$  is time in years and  $h$  is height in inches. The seedlings are 6 inches tall when planted ( $t = 0$ ).

- (a) Find the height function.  
 (b) How tall are the shrubs when they are sold?

56. **Cash Flow** The rate of disbursement  $dQ/dt$  of a \$2 million federal grant is proportional to the square of  $100 - t$ , where  $t$  is the time (in days,  $0 \leq t \leq 100$ ) and  $Q$  is the amount that remains to be disbursed. Find the amount that remains to be disbursed after 50 days. Assume that the entire grant will be disbursed after 100 days.

**Marginal Propensity to Consume** In Exercises 57 and 58, (a) use the marginal propensity to consume,  $dQ/dx$ , to write  $Q$  as a function of  $x$ , where  $x$  is the income (in dollars) and  $Q$  is the income consumed (in dollars). Assume that 100% of the income is consumed for families that have annual incomes of \$20,000 or less. (b) Use the result of part (a) to complete the table showing the income consumed and the income saved,  $x - Q$ , for various incomes. (c) Use a graphing utility to represent graphically the income consumed and saved.

$x$	20,000	50,000	100,000	150,000
$Q$				
$x - Q$				

57.  $\frac{dQ}{dx} = \frac{0.95}{(x - 19,999)^{0.05}}$ ,  $x \geq 20,000$

58.  $\frac{dQ}{dx} = \frac{0.93}{(x - 19,999)^{0.07}}$ ,  $x \geq 20,000$

In Exercises 59 and 60, use a symbolic integration utility to find the indefinite integral. Verify the result by differentiating.

59.  $\int \frac{1}{\sqrt{x} + \sqrt{x + 1}} dx$       60.  $\int \frac{x}{\sqrt{3x + 2}} dx$

Using th

Each of th  
integration

Integra

Let u b

EXAM

Find each

(a)  $\int 2e^x dx$

(b)  $\int 2e^{2x} dx$

(c)  $\int e^{x^2} dx$

SOLUTION

(a)  $\int 2e^x dx$

(b)  $\int 2e^{2x} dx$

(c)  $\int e^{x^2} dx$

You can cl

**PREREQUISITE  
REVIEW 5.3**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1 and 2, find the domain of the function.

1.  $y = \ln(2x - 5)$

2.  $y = \ln(x^2 - 5x + 6)$

In Exercises 3–6, use long division to rewrite the quotient.

3.  $\frac{x^2 + 4x + 2}{x + 2}$

4.  $\frac{x^2 - 6x + 9}{x - 4}$

5.  $\frac{x^3 + 4x^2 - 30x - 4}{x^2 - 4x}$

6.  $\frac{x^4 - x^3 + x^2 + 15x + 2}{x^2 + 5}$

In Exercises 7–10, evaluate the integral.

7.  $\int \left( x^3 + \frac{1}{x^2} \right) dx$

8.  $\int \frac{x^2 + 2x}{x} dx$

9.  $\int \frac{x^3 + 4}{x^2} dx$

10.  $\int \frac{x + 3}{x^3} dx$

**EXERCISES 5.3**

In Exercises 1–12, use the Exponential Rule to find the indefinite integral.

1.  $\int 2e^{2x} dx$

2.  $\int -3e^{-3x} dx$

3.  $\int e^{4x} dx$

4.  $\int e^{-0.25x} dx$

5.  $\int 9xe^{-x^2} dx$

6.  $\int 3xe^{0.5x^2} dx$

7.  $\int 5x^2 e^{x^3} dx$

8.  $\int (2x + 1)e^{x^2+x} dx$

9.  $\int (x^2 + 2x)e^{x^3+3x^2-1} dx$

10.  $\int 3(x - 4)e^{x^2-8x} dx$

11.  $\int 5e^{2-x} dx$

12.  $\int 3e^{-(x+1)} dx$

In Exercises 13–28, use the Log Rule to find the indefinite integral.

13.  $\int \frac{1}{x+1} dx$

14.  $\int \frac{1}{x-5} dx$

15.  $\int \frac{1}{3-2x} dx$

16.  $\int \frac{1}{6x-5} dx$

17.  $\int \frac{2}{3x+5} dx$

18.  $\int \frac{5}{2x-1} dx$

19.  $\int \frac{x}{x^2+1} dx$

20.  $\int \frac{x^2}{3-x^3} dx$

21.  $\int \frac{x^2}{x^3+1} dx$

22.  $\int \frac{x}{x^2+4} dx$

23.  $\int \frac{x+3}{x^2+6x+7} dx$

24.  $\int \frac{x^2+2x+3}{x^3+3x^2+9x+1} dx$

25.  $\int \frac{1}{x \ln x} dx$

26.  $\int \frac{1}{x(\ln x)^2} dx$

27.  $\int \frac{e^{-x}}{1-e^{-x}} dx$

28.  $\int \frac{e^x}{1+e^x} dx$

 In Exercises 29–38, use a symbolic integration utility to find the indefinite integral.

29.  $\int \frac{1}{x^2} e^{2/x} dx$

30.  $\int \frac{1}{x^3} e^{1/4x^2} dx$

31.  $\int \frac{1}{\sqrt{x}} e^{\sqrt{x}} dx$

32.  $\int \frac{e^{1/\sqrt{x}}}{x^{3/2}} dx$

33.  $\int (e^x - 2)^2 dx$

34.  $\int (e^x - e^{-x})^2 dx$

35.  $\int \frac{e^{-x}}{1+e^{-x}} dx$

36.  $\int \frac{3e^x}{2+e^x} dx$

37.  $\int \frac{4e^{2x}}{5-e^{2x}} dx$

38.  $\int \frac{-e^{3x}}{2-e^{3x}} dx$

In Exercises 39–54  
find the indefin

39.  $\int \frac{e^{2x} + 2e^x}{e^x} dx$

41.  $\int e^x \sqrt{1-e^x} dx$

43.  $\int \frac{1}{(x-1)^2} dx$

45.  $\int 4e^{2x-1} dx$

47.  $\int \frac{x^3 - 8x}{2x^2} dx$

49.  $\int \frac{2}{1+e^{-x}} dx$

51.  $\int \frac{x^2 + 2x}{x-1} dx$

53.  $\int \frac{1+e^{-x}}{1+xe^{-x}} dx$

In Exercises 55 and  
graph passes thr

55.  $f'(x) = \frac{x^2}{1-x^3}$

56.  $f'(x) = \frac{x^3}{1-x^3}$

57. **Biology** A

$$\frac{dP}{dt} = \frac{P}{1-t}$$

where  $t$  is  $t$   
1000.

(a) Write  
terms  $C$

(b) What  $i$

(c) After  $h$

58. **Biology**

trout popul  
change  $C$

$$\frac{dP}{dt} = \frac{P}{1-t}$$

where  $t$  is  
2500.

(a) Write  
terms

(b) What

(c) Accor  
entire

In Exercises 39–54, use any basic integration formula or formulas to find the indefinite integral.

39.  $\int \frac{e^{2x} + 2e^x + 1}{e^x} dx$

40.  $\int (6x + e^x) \sqrt{3x^2 + e^x} dx$

41.  $\int e^x \sqrt{1 - e^x} dx$

42.  $\int \frac{2(e^x - e^{-x})}{(e^x + e^{-x})^2} dx$

43.  $\int \frac{1}{(x-1)^2} dx$

44.  $\int \frac{1}{\sqrt{x+1}} dx$

45.  $\int 4e^{2x-1} dx$

46.  $\int (5e^{-2x} + 1) dx$

47.  $\int \frac{x^3 - 8x}{2x^2} dx$

48.  $\int \frac{x-1}{4x} dx$

49.  $\int \frac{2}{1 + e^{-x}} dx$

50.  $\int \frac{3}{1 + e^{-3x}} dx$

51.  $\int \frac{x^2 + 2x + 5}{x-1} dx$

52.  $\int \frac{x-3}{x+3} dx$

53.  $\int \frac{1 + e^{-x}}{1 + xe^{-x}} dx$

54.  $\int \frac{5}{e^{-5x} + 7} dx$

In Exercises 55 and 56, find the equation of the function  $f$  whose graph passes through the point.

55.  $f'(x) = \frac{x^2 + 4x + 3}{x-1}$ ; (2, 4)

56.  $f'(x) = \frac{x^3 - 4x^2 + 3}{x-3}$ ; (4, -1)

57. **Biology** A population of bacteria is growing at the rate of

$$\frac{dP}{dt} = \frac{3000}{1 + 0.25t}$$

where  $t$  is the time in days. When  $t = 0$ , the population is 1000.

(a) Write an equation that models the population  $P$  in terms of the time  $t$ .

(b) What is the population after 3 days?

(c) After how many days will the population be 12,000?

58. **Biology** Because of an insufficient oxygen supply, the trout population in a lake is dying. The population's rate of change can be modeled by

$$\frac{dP}{dt} = -125e^{-t/20}$$

where  $t$  is the time in days. When  $t = 0$ , the population is 2500.

(a) Write an equation that models the population  $P$  in terms of the time  $t$ .

(b) What is the population after 15 days?

(c) According to this model, how long will it take for the entire trout population to die?

59. **Demand** The marginal price for the demand of a product can be modeled by  $dp/dx = 0.1e^{-x/500}$ , where  $x$  is the quantity demanded. When the demand is 600 units, the price is \$30.

(a) Find the demand function,  $p = f(x)$ .

(b) Use a graphing utility to graph the demand function. Does price increase or decrease as demand increases?

(c) Use the *zoom* and *trace* features of the graphing utility to find the quantity demanded when the price is \$22.

60. **Revenue** The marginal revenue for the sale of a product can be modeled by

$$\frac{dR}{dx} = 50 - 0.02x + \frac{100}{x+1}$$

where  $x$  is the quantity demanded.

(a) Find the revenue function.

(b) Use a graphing utility to graph the revenue function.

(c) Find the revenue when 1500 units are sold.

(d) Use the *zoom* and *trace* features of the graphing utility to find the number of units sold when the revenue is \$60,230.

61. **Average Salary** From 1995 through 2002, the average salary for superintendents  $S$  (in dollars) in the United States changed at the rate of

$$\frac{dS}{dt} = 2621.7e^{0.07t}$$

where  $t = 5$  corresponds to 1995. In 2001, the average salary for superintendents was \$118,496. (Source: Educational Research Service)

(a) Write a model that gives the average salary for superintendents per year.

(b) Use the model to find the average salary for superintendents in 1999.

62. **Sales** The rate of change in sales for The Yankee Candle Company from 1998 through 2003 can be modeled by

$$\frac{dS}{dt} = 1.04t + \frac{544.694}{t}$$

where  $S$  is the sales (in millions) and  $t = 8$  corresponds to 1998. In 1999, the sales for The Yankee Candle Company were \$256.6 million. (Source: The Yankee Candle Company)

(a) Find a model for sales from 1998 through 2003.

(b) Find The Yankee Candle Company's sales in 2002.

**True or False?** In Exercises 63 and 64, determine whether the statement is true or false. If it is false, explain why or give an example that shows it is false.

63.  $(\ln x)^{1/2} = \frac{1}{2}(\ln x)$

64.  $\int \ln x = \left(\frac{1}{x}\right) + C$

**PREREQUISITE  
REVIEW 5.4**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–4, find the indefinite integral.

1.  $\int (3x + 7) dx$

2.  $\int (x^{3/2} + 2\sqrt{x}) dx$

3.  $\int \frac{1}{5x} dx$

4.  $\int e^{-6x} dx$

In Exercises 5 and 6, evaluate the expression when  $a = 5$  and  $b = 3$ .

5.  $\left(\frac{a}{5} - a\right) - \left(\frac{b}{5} - b\right)$

6.  $\left(6a - \frac{a^3}{3}\right) - \left(6b - \frac{b^3}{3}\right)$

In Exercises 7–10, integrate the marginal function.

7.  $\frac{dC}{dx} = 0.02x^{3/2} + 29,500$

8.  $\frac{dR}{dx} = 9000 + 2x$

9.  $\frac{dP}{dx} = 25,000 - 0.01x$

10.  $\frac{dC}{dx} = 0.03x^2 + 4600$

**EXERCISES 5.4**

In Exercises 1–8, sketch the region whose area is represented by the definite integral. Then use a geometric formula to evaluate the integral.

1.  $\int_0^2 3 dx$

2.  $\int_0^4 2 dx$

3.  $\int_0^5 (x + 1) dx$

4.  $\int_0^3 (2x + 1) dx$

5.  $\int_{-2}^3 |x - 1| dx$

6.  $\int_{-1}^4 |x - 2| dx$

7.  $\int_{-3}^3 \sqrt{9 - x^2} dx$

8.  $\int_0^2 \sqrt{4 - x^2} dx$

In Exercises 9 and 10, use the values  $\int_0^5 f(x) dx = 8$  and  $\int_0^5 g(x) dx = 3$  to evaluate the definite integral.

9. (a)  $\int_0^5 [f(x) + g(x)] dx$

(b)  $\int_0^5 [f(x) - g(x)] dx$

(c)  $\int_0^5 -4f(x) dx$

(d)  $\int_0^5 [f(x) - 3g(x)] dx$

10. (a)  $\int_0^5 2g(x) dx$

(b)  $\int_5^0 f(x) dx$



(c)  $\int_5^0 f(x) dx$

(d)  $\int_0^5 [f(x) - f(x)] dx$

In Exercises 11–18, find the area of the region.



11.  $y = x - x^2$

12.  $y = 1 - x^4$





13.  $y = \frac{1}{x^2}$

14.  $y = \frac{2}{\sqrt{x}}$




15.  $y = 3e^{-x/2}$

16.  $y = 2e^{x/2}$






17.  $y = \frac{x^2 + 4}{x}$



18.  $y = \frac{x - 2}{x}$



In Exercises 19–42, evaluate the definite integral.

- |   |   |
|---|---|
| 19. $\int_0^1 2x \, dx$                       | 20. $\int_2^7 3v \, dv$                               |
| 21. $\int_{-1}^0 (2x + 1) \, dx$              | 22. $\int_2^5 (-3x + 4) \, dx$                        |
| 23. $\int_{-1}^1 (2t - 1)^2 \, dt$            | 24. $\int_0^1 (1 - 2x)^2 \, dx$                       |
| 25. $\int_0^3 (x - 2)^3 \, dx$                | 26. $\int_2^4 (x - 3)^4 \, dx$                        |
| 27. $\int_{-1}^1 (\sqrt[3]{t} - 2) \, dt$     | 28. $\int_1^4 \sqrt{\frac{2}{x}} \, dx$               |
| 29. $\int_1^4 \frac{2u - 1}{\sqrt{u}} \, du$  | 30. $\int_0^1 \frac{x - \sqrt{x}}{3} \, dx$           |
| 31. $\int_{-1}^0 (t^{1/3} - t^{2/3}) \, dt$   | 32. $\int_0^4 (x^{1/2} + x^{1/4}) \, dx$              |
| 33. $\int_0^4 \frac{1}{\sqrt{2x + 1}} \, dx$  | 34. $\int_0^2 \frac{x}{\sqrt{1 + 2x^2}} \, dx$        |
| 35. $\int_0^1 e^{-2x} \, dx$                  | 36. $\int_1^2 e^{1-x} \, dx$                          |
| 37. $\int_1^3 \frac{e^{3/x}}{x^2} \, dx$      | 38. $\int_{-1}^1 (e^x - e^{-x}) \, dx$                |
| 39. $\int_0^1 e^{2x} \sqrt{e^{2x} + 1} \, dx$ | 40. $\int_0^1 \frac{e^{-x}}{\sqrt{e^{-x} + 1}} \, dx$ |
| 41. $\int_0^2 \frac{x}{1 + 4x^2} \, dx$       | 42. $\int_0^1 \frac{e^{2x}}{e^{2x} + 1} \, dx$        |

In Exercises 43–46, evaluate the definite integral by the most convenient method. Explain your approach.

- |                                    |                                   |
|------------------------------------|-----------------------------------|
| 43. $\int_{-1}^1  4x  \, dx$       | 44. $\int_0^3  2x - 3  \, dx$     |
| 45. $\int_0^4 (2 -  x - 2 ) \, dx$ | 46. $\int_{-4}^4 (4 -  x ) \, dx$ |

In Exercises 47–50, evaluate the definite integral by hand. Then use a symbolic integration utility to evaluate the definite integral. Briefly explain any differences in your results.

- |   |   |
|---|---|
| 47. $\int_{-1}^2 \frac{x}{x^2 - 9} \, dx$ | 48. $\int_2^3 \frac{x + 1}{x^2 + 2x - 3} \, dx$ |
| 49. $\int_0^3 \frac{2e^x}{2 + e^x} \, dx$ | 50. $\int_1^2 \frac{(2 + \ln x)^3}{x} \, dx$    |

In Exercises 51–56, evaluate the definite integral by hand. Then use a graphing utility to graph the region whose area is represented by the integral.

- |   |  |
|---|--|
| 51. $\int_1^3 (4x - 3) \, dx$             | 52. $\int_0^2 (x + 4) \, dx$             |
| 53. $\int_0^1 (x - x^3) \, dx$            | 54. $\int_0^1 \sqrt{x}(1 - x) \, dx$     |
| 55. $\int_2^4 \frac{3x^2}{x^3 - 1} \, dx$ | 56. $\int_0^{\ln 6} \frac{e^x}{2} \, dx$ |

In Exercises 57–60, find the area of the region bounded by the graphs. Use a graphing utility to graph the region and verify your results.

57.  $y = 3x^2 + 1$ ,  $y = 0$ ,  $x = 0$ , and  $x = 2$   
 58.  $y = 1 + \sqrt{x}$ ,  $y = 0$ ,  $x = 0$ , and  $x = 4$   
 59.  $y = (x + 5)/x$ ,  $y = 0$ ,  $x = 1$ , and  $x = 5$   
 60.  $y = 3e^x$ ,  $y = 0$ ,  $x = -2$ , and  $x = 1$

In Exercises 61–68, use a graphing utility to graph the function over the interval. Find the average value of the function over the interval. Then find all  $x$ -values in the interval for which the function is equal to its average value.

- | Function                         | Interval  |
|----------------------------------|-----------|
| 61. $f(x) = 6 - x^2$             | $[-2, 2]$ |
| 62. $f(x) = x - 2\sqrt{x}$       | $[0, 4]$  |
| 63. $f(x) = 5e^{0.2(x-10)}$      | $[0, 10]$ |
| 64. $f(x) = 2e^{x/4}$            | $[0, 4]$  |
| 65. $f(x) = x\sqrt{4 - x^2}$     | $[0, 2]$  |
| 66. $f(x) = \frac{1}{(x - 3)^2}$ | $[0, 2]$  |
| 67. $f(x) = \frac{5x}{x^2 + 1}$  | $[0, 7]$  |
| 68. $f(x) = \frac{4x}{x^2 + 1}$  | $[0, 1]$  |

In Exercises 69–72, state whether the function is even, odd, or neither.

- |                          |                       |
|--------------------------|-----------------------|
| 69. $f(x) = 3x^4$        | 70. $g(x) = x^3 - 2x$ |
| 71. $g(t) = 2t^5 - 3t^2$ | 72. $f(t) = 5t^4 + 1$ |

73. Use the value  $\int_0^2 x^2 \, dx = \frac{8}{3}$  to evaluate each definite integral. Explain your reasoning.

- |                             |                             |                           |
|-----------------------------|-----------------------------|---------------------------|
| (a) $\int_{-2}^0 x^2 \, dx$ | (b) $\int_{-2}^2 x^2 \, dx$ | (c) $\int_0^2 -x^2 \, dx$ |
|-----------------------------|-----------------------------|---------------------------|

74. Use the value  $\int_0^2 x^3 \, dx = 4$  to evaluate each definite integral. Explain your reasoning.

- |                             |                             |                           |
|-----------------------------|-----------------------------|---------------------------|
| (a) $\int_{-2}^0 x^3 \, dx$ | (b) $\int_{-2}^2 x^3 \, dx$ | (c) $\int_0^2 3x^3 \, dx$ |
|-----------------------------|-----------------------------|---------------------------|

**Marginal A** in cost  $C$ , revenue  $R$ , and profit  $P$  are functions of quantity  $q$ . In each case, assume the functions are specified values.

- Marginal**
75.  $\frac{dC}{dq} = 2.2$   
 76.  $\frac{dC}{dq} = \frac{20}{q}$   
 77.  $\frac{dR}{dq} = 48$   
 78.  $\frac{dR}{dq} = 75\sqrt{q}$   
 79.  $\frac{dP}{dq} = \frac{400}{q}$   
 80.  $\frac{dP}{dq} = 12$

**Annuity** In each case, assume the annuity is paid monthly with income  $f(t)$  dollars per month.

81.  $c(t) = \$2$   
 82.  $c(t) = \$5$   
 83.  $c(t) = \$1$   
 84.  $c(t) = \$2$

**Capital Accumulation** In each case, assume the rate of investment is  $f(t)$  dollars per year over a five-year period.

**Capital Accumulation**

where  $t$  is the time in years.

85.  $\frac{dI}{dt} = 500$   
 86.  $\frac{dI}{dt} = 100$   
 87.  $\frac{dI}{dt} = 500$   
 88.  $\frac{dI}{dt} = \frac{12}{t^2}$

89. **Cost** The total cost of equipment is  $C(t)$  dollars after  $t$  years.

$C = 5$

Find the total cost after 5 years.

**Marginal Analysis** In Exercises 75–80, find the change in cost  $C$ , revenue  $R$ , or profit  $P$ , for the given marginal. In each case, assume that the number of units  $x$  increases by 3 from the specified value of  $x$ .

	Marginal	Number of Units, $x$
75.	$\frac{dC}{dx} = 2.25$	$x = 100$
76.	$\frac{dC}{dx} = \frac{20,000}{x^2}$	$x = 10$
77.	$\frac{dR}{dx} = 48 - 3x$	$x = 12$
78.	$\frac{dR}{dx} = 75\left(20 + \frac{900}{x}\right)$	$x = 500$
79.	$\frac{dP}{dx} = \frac{400 - x}{150}$	$x = 200$
80.	$\frac{dP}{dx} = 12.5(40 - 3\sqrt{x})$	$x = 125$

**Annuity** In Exercises 81–84, find the amount of an annuity with income function  $c(t)$ , interest rate  $r$ , and term  $T$ .

81.  $c(t) = \$250$ ,  $r = 8\%$ ,  $T = 6$  years  
 82.  $c(t) = \$500$ ,  $r = 9\%$ ,  $T = 4$  years  
 83.  $c(t) = \$1500$ ,  $r = 2\%$ ,  $T = 10$  years  
 84.  $c(t) = \$2000$ ,  $r = 3\%$ ,  $T = 15$  years

**Capital Accumulation** In Exercises 85–88, you are given the rate of investment  $dl/dt$ . Find the capital accumulation over a five-year period by evaluating the definite integral

$$\text{Capital accumulation} = \int_0^5 \frac{dl}{dt} dt$$

where  $t$  is the time in years.

85.  $\frac{dl}{dt} = 500$   
 86.  $\frac{dl}{dt} = 100t$   
 87.  $\frac{dl}{dt} = 500\sqrt{t+1}$   
 88.  $\frac{dl}{dt} = \frac{12,000t}{(t^2+2)^2}$

89. **Cost** The total cost of purchasing and maintaining a piece of equipment for  $x$  years can be modeled by

$$C = 5000\left(25 + 3 \int_0^x t^{1/4} dt\right).$$

Find the total cost after (a) 1 year, (b) 5 years, and (c) 10 years.

90. **Depreciation** A company purchases a new machine for which the rate of depreciation can be modeled by

$$\frac{dV}{dt} = 10,000(t - 6), \quad 0 \leq t \leq 5$$

where  $V$  is the value of the machine after  $t$  years. Set up and evaluate the definite integral that yields the total loss of value of the machine over the first 3 years.

91. **Compound Interest** A deposit of \$2250 is made in a savings account at an annual interest rate of 12%, compounded continuously. Find the average balance in the account during the first 5 years.

92. **Mortgage Debt** The rate of change of mortgage debt outstanding for one- to four-family homes in the United States from 1993 through 2002 can be modeled by

$$\frac{dM}{dt} = 5.4399t^2 + 6603.7e^{-t}$$

where  $M$  is the mortgage debt outstanding (in billions of dollars) and  $t = 3$  corresponds to 1993. In 1993, the mortgage debt outstanding in the United States was \$3119 billion. (Source: Board of Governors of the Federal Reserve System)

- (a) Write a model for the debt as a function of  $t$ .  
 (b) What was the average mortgage debt outstanding for 1993 through 2002?

93. **Medicine** The velocity  $v$  of blood at a distance  $r$  from the center of an artery of radius  $R$  can be modeled by

$$v = k(R^2 - r^2)$$

where  $k$  is a constant. Find the average velocity along a radius of the artery. (Use 0 and  $R$  as the limits of integration.)

94. **Biology** The rate of change in the number of coyotes  $N(t)$  in a population is directly proportional to  $650 - N(t)$ , where  $t$  is time in years.

$$\frac{dN}{dt} = k[650 - N(t)]$$

When  $t = 0$ , the population is 300, and when  $t = 2$ , the population has increased to 500.

- (a) Find the population function.  
 (b) Find the average number of coyotes over the first 5 years.

⊕ In Exercises 95–98, use a symbolic integration utility to evaluate the definite integral.

95.  $\int_3^6 \frac{x}{3\sqrt{x^2-8}} dx$

96.  $\int_{1/2}^1 (x+1)\sqrt{1-x} dx$

97.  $\int_2^5 \left(\frac{1}{x^2} - \frac{1}{x^3}\right) dx$

98.  $\int_0^1 x^3(x^3+1)^3 dx$

**PREREQUISITE REVIEW 5.5**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–4, simplify the expression.

1.  $(-x^2 + 4x + 3) - (x + 1)$
2.  $(-2x^2 + 3x + 9) - (-x + 5)$
3.  $(-x^3 + 3x^2 - 1) - (x^2 - 4x + 4)$
4.  $(3x + 1) - (-x^3 + 9x + 2)$


In Exercises 5–10, find the points of intersection of the graphs.

5.  $f(x) = x^2 - 4x + 4$ ,  $g(x) = 4$
6.  $f(x) = -3x^2$ ,  $g(x) = 6 - 9x$
7.  $f(x) = x^2$ ,  $g(x) = -x + 6$
8.  $f(x) = \frac{1}{2}x^3$ ,  $g(x) = 2x$
9.  $f(x) = x^2 - 3x$ ,  $g(x) = 3x - 5$
10.  $f(x) = e^x$ ,  $g(x) = e$


**EXERCISES 5.5**

In Exercises 1–8, find the area of the region.


1.  $f(x) = x^2 - 6x$   
 $g(x) = 0$




2.  $f(x) = x^2 + 2x + 1$   
 $g(x) = 2x + 5$




3.  $f(x) = x^2 - 4x + 3$   
 $g(x) = -x^2 + 2x + 3$




4.  $f(x) = x^2$   
 $g(x) = x^3$




5.  $f(x) = 3(x^3 - x)$   
 $g(x) = 0$




6.  $f(x) = (x - 1)^3$   
 $g(x) = x - 1$



7.  $f(x) = e^x - 1$   
 $g(x) = 0$



8.  $f(x) = -x + 3$   
 $g(x) = 2x^{-1}$



In Exercises 9–14, find the definite integral.

9.  $\int_0^4 [(x + 1) - 2] dx$
10.  $\int_{-1}^1 [(1 - x) - 2] dx$
11.  $\int_{-2}^2 [2x^2 - 1] dx$
12.  $\int_{-4}^0 [(x - 1) - 2] dx$
13.  $\int_{-1}^2 [(y^2 - 1) - 2] dy$
14.  $\int_{-2}^3 [(y - 1) - 2] dy$

In Exercises 15–26, find the area of the region.

15.  $y = \frac{1}{x^2}$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
16.  $y = x^3 - 1$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
17.  $f(x) = x^2 - 1$ ,  $g(x) = x - 1$
18.  $f(x) = x^2 - 1$ ,  $g(x) = x - 1$
19.  $y = x^2 - 1$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
20.  $y = 4 - x^2$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
21.  $y = xe^{-x}$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
22.  $y = \frac{e^{1/x}}{x^2}$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
23.  $y = \frac{8}{x^3}$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
24.  $y = \frac{1}{x^2}$ ,  $x = 1$ ,  $x = 2$ ,  $y = 0$
25.  $f(x) = e^x - 1$ ,  $g(x) = x - 1$
26.  $f(x) = \frac{1}{x^2}$ ,  $g(x) = x - 1$

In Exercises 27–30, find the area of the region.

27.  $f(y) = y^2 - 1$ ,  $g(y) = y - 1$
28.  $f(y) = y^2 - 1$ ,  $g(y) = y - 1$
29.  $f(y) = y^2 - 1$ ,  $g(y) = y - 1$
30.  $f(y) = y^2 - 1$ ,  $g(y) = y - 1$

In Exercises 9–14, sketch the region whose area is represented by the definite integral.

9.  $\int_0^4 [(x+1) - \frac{1}{2}x] dx$

10.  $\int_{-1}^1 [(1-x^2) - (x^2-1)] dx$

11.  $\int_{-2}^2 [2x^2 - (x^4 - 2x^2)] dx$

12.  $\int_{-4}^0 [(x-6) - (x^2 + 5x - 6)] dx$

13.  $\int_{-1}^2 [(y^2 + 2) - 1] dy$

14.  $\int_{-2}^3 [(y+6) - y^2] dy$

In Exercises 15–26, sketch the region bounded by the graphs of the functions and find the area of the region.

15.  $y = \frac{1}{x^2}, y = 0, x = 1, x = 5$

16.  $y = x^3 - 2x + 1, y = -2x, x = 1$

17.  $f(x) = \sqrt[3]{x}, g(x) = x$

18.  $f(x) = \sqrt{3x+1}, g(x) = x+1$

19.  $y = x^2 - 4x + 3, y = 3 + 4x - x^2$

20.  $y = 4 - x^2, y = x^2$

21.  $y = xe^{-x^2}, y = 0, x = 0, x = 1$

22.  $y = \frac{e^{1/x}}{x^2}, y = 0, x = 1, x = 3$

23.  $y = \frac{8}{x}, y = x^2, y = 0, x = 1, x = 4$

24.  $y = \frac{1}{x}, y = x^3, x = \frac{1}{2}, x = 1$

25.  $f(x) = e^{0.5x}, g(x) = -\frac{1}{x}, x = 1, x = 2$

26.  $f(x) = \frac{1}{x}, g(x) = -e^x, x = \frac{1}{2}, x = 1$

In Exercises 27–30, sketch the region bounded by the graphs of the functions and find the area of the region.

27.  $f(y) = y^2, g(y) = y + 2$

28.  $f(y) = y(2-y), g(y) = -y$

29.  $f(y) = \sqrt{y}, y = 9, x = 0$

30.  $f(y) = y^2 + 1, g(y) = 4 - 2y$

In Exercises 31–34, use a graphing utility to graph the region bounded by the graphs of the functions. Write the definite integrals that represent the area of the region. (Hint: Multiple integrals may be necessary.)

31.  $f(x) = 2x, g(x) = 4 - 2x, h(x) = 0$

32.  $f(x) = x(x^2 - 3x + 3), g(x) = x^2$

33.  $y = \frac{4}{x}, y = x, x = 1, x = 4$

34.  $y = x^3 - 4x^2 + 1, y = x - 3$

In Exercises 35–38, use a graphing utility to graph the region bounded by the graphs of the functions, and find the area of the region.

35.  $f(x) = x^2 - 4x, g(x) = 0$

36.  $f(x) = 3 - 2x - x^2, g(x) = 0$

37.  $f(x) = x^2 + 2x + 1, g(x) = x + 1$

38.  $f(x) = -x^2 + 4x + 2, g(x) = x + 2$

In Exercises 39 and 40, use integration to find the area of the triangular region having the given vertices.

39. (0, 0), (4, 0), (4, 4)

40. (0, 0), (4, 0), (6, 4)

**Consumer and Producer Surpluses** In Exercises 41–46, find the consumer and producer surpluses.

*Demand Function*

*Supply Function*

41.  $p_1(x) = 50 - 0.5x$

$p_2(x) = 0.125x$

42.  $p_1(x) = 300 - x$

$p_2(x) = 100 + x$

43.  $p_1(x) = 200 - 0.02x^2$

$p_2(x) = 100 + x$

44.  $p_1(x) = 1000 - 0.4x^2$

$p_2(x) = 42x$

45.  $p_1(x) = \frac{10,000}{\sqrt{x+100}}$

$p_2(x) = 100\sqrt{0.05x+10}$

46.  $p_1(x) = \sqrt{25 - 0.1x}$

$p_2(x) = \sqrt{9 + 0.1x} - 2$

47. **Writing** Describe the characteristics of typical demand and supply functions.

48. **Writing** Suppose that the demand and supply functions for a product do not intersect. What can you conclude?

**Revenue** In Exercises 49 and 50, two models,  $R_1$  and  $R_2$ , are given for revenue (in billions of dollars per year) for a large corporation. Both models are estimates of revenues for 2004–2008, with  $t = 4$  corresponding to 2004. Which model is projecting the greater revenue? How much more total revenue does that model project over the four-year period?

49.  $R_1 = 7.21 + 0.58t, R_2 = 7.21 + 0.45t$

50.  $R_1 = 7.21 + 0.26t + 0.02t^2, R_2 = 7.21 + 0.1t + 0.01t^2$

**51. Fuel Cost** The projected fuel cost  $C$  (in millions of dollars per year) for an airline company from 2004 through 2010 is  $C_1 = 568.5 + 7.15t$ , where  $t = 4$  corresponds to 2004. If the company purchases more efficient airplane engines, fuel cost is expected to decrease and to follow the model  $C_2 = 525.6 + 6.43t$ . How much can the company save with the more efficient engines? Explain your reasoning.

**52. Health** An epidemic was spreading such that  $t$  weeks after its outbreak it had infected

$$N_1(t) = 0.1t^2 + 0.5t + 150, \quad 0 \leq t \leq 50$$

people. Twenty-five weeks after the outbreak, a vaccine was developed and administered to the public. At that point, the number of people infected was governed by the model

$$N_2(t) = -0.2t^2 + 6t + 200.$$

Approximate the number of people that the vaccine prevented from becoming ill during the epidemic.

**53. Consumer Trends** For the years 1990 through 2001, the per capita consumption of tomatoes (in pounds per year) in the United States can be modeled by

$$C(t) = \begin{cases} 0.085t^3 - 0.309t^2 + 0.13t + 15.5, & 0 \leq t \leq 4 \\ 0.01515t^4 - 0.5348t^3 + 6.864t^2 - 37.68t + 91.4 & 4 < t \leq 11 \end{cases}$$

where  $t = 0$  corresponds to 1990. (Source: U.S. Department of Agriculture)

- (a) Use a graphing utility to graph this model.
- (b) Suppose the tomato consumption from 1995 through 2001 had continued to follow the model for 1990 through 1994. How many more or fewer pounds of tomatoes would have been consumed from 1995 through 2001?

**54. Consumer and Producer Surpluses** Factory orders for an air conditioner are about 6000 units per week when the price is \$331 and about 8000 units per week when the price is \$303. The supply function is given by  $p = 0.0275x$ . Find the consumer and producer surpluses. (Assume the demand function is linear.)

**55. Consumer and Producer Surpluses** Repeat Exercise 54 with a demand of about 6000 units per week when the price is \$325 and about 8000 units per week when the price is \$300. Find the consumer and producer surpluses. (Assume the demand function is linear.)

**56. Cost, Revenue, and Profit** The revenue from a manufacturing process (in millions of dollars per year) is projected to follow the model  $R = 100$  for 10 years. Over the same period of time, the cost (in millions of dollars per year) is projected to follow the model  $C = 60 + 0.2t^2$ , where  $t$  is the time (in years). Approximate the profit over the 10-year period.

**57. Cost, Revenue, and Profit** Repeat Exercise 56 for revenue and cost models given by  $R = 100 + 0.08t$  and  $C = 60 + 0.2t^2$ .

**58. Lorenz Curve** Economists use *Lorenz curves* to illustrate the distribution of income in a country. Letting  $x$  represent the percent of families in a country and  $y$  the percent of total income, the model  $y = x$  would represent a country in which each family had the same income. The Lorenz curve,  $y = f(x)$ , represents the actual income distribution. The area between these two models, for  $0 \leq x \leq 100$ , indicates the “income inequality” of a country. In 2001, the Lorenz curve for the United States could be modeled by

$$y = (0.00059x^2 + 0.0233x + 1.731)^2, \quad 0 \leq x \leq 100$$

where  $x$  is measured from the poorest to the wealthiest families. Find the income inequality for the United States in 2001. (Source: U.S. Census Bureau)

**59. Income Distribution** Using the Lorenz curve in Exercise 58, complete the table, which lists the percent of total income earned by each quintile in the United States in 2001.

Quintile	Lowest	2nd	3rd	4th	Highest
Percent					

### BUSINESS CAPSULE



In 1994, wardrobe consultant and personal shopper Marilyn N. Wright started the company Marilyn's Fashions in Newark, Delaware. Wright consults with her 450 American and 250 international clients on their wardrobes, styles, and budgets, then delivers the clothes and accessories to their doorsteps in less than 2 weeks. She used \$5000 in start-up capital while working as a claims adjuster and now brings in \$250,000 in annual revenue.

**60. Research Project** Use your school's library, the Internet, or some other reference source to research a small company similar to that described above. Describe the impact of different factors, such as start-up capital and market conditions, on a company's revenue.

### The Mid

In Section Calculus to integrand. I the integral Midpoint F

### EXAM

Use the five bounded by  $x = 2$ .

**SOLUTION** midpoint of

$$\left[0, \frac{2}{5}\right]$$

The width c

Area  $\approx$

### TRY IT

Use four r graph of  $f$

For the gral. That is

Area =

**PREREQUISITE REVIEW 5.6**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, find the midpoint of the interval.

1.  $[0, \frac{1}{3}]$

3.  $[\frac{3}{20}, \frac{4}{20}]$

5.  $[2, \frac{31}{15}]$

2.  $[\frac{1}{10}, \frac{2}{10}]$

4.  $[1, \frac{7}{6}]$

6.  $[\frac{26}{9}, 3]$

In Exercises 7–10, find the limit.

7.  $\lim_{x \rightarrow \infty} \frac{2x^2 + 4x - 1}{3x^2 - 2x}$

8.  $\lim_{x \rightarrow \infty} \frac{4x + 5}{7x - 5}$



9.  $\lim_{x \rightarrow \infty} \frac{x - 7}{x^2 + 1}$

10.  $\lim_{x \rightarrow \infty} \frac{5x^3 + 1}{x^3 + x^2 + 4}$


**EXERCISES 5.6**

In Exercises 1–4, use the Midpoint Rule with  $n = 4$  to approximate the area of the region. Compare your result with the exact area obtained with a definite integral.


1.  $f(x) = -2x + 3, [0, 1]$     2.  $f(x) = \frac{1}{x}, [1, 5]$



3.  $f(x) = \sqrt{x}, [0, 1]$



4.  $f(x) = 1 - x^2, [-1, 1]$



In Exercises 5–12, use the Midpoint Rule with  $n = 4$  to approximate the area of the region bounded by the graph of  $f$  and the  $x$ -axis over the interval. Compare your result with the exact area. Sketch the region.

Function	Interval
5. $f(x) = x^2 + 2$	$[-1, 1]$
6. $f(x) = 4 - x^2$	$[0, 2]$
7. $f(x) = 2x^2$	$[1, 3]$
8. $f(x) = 2x - x^3$	$[0, 1]$
9. $f(x) = x^2 - x^3$	$[0, 1]$
10. $f(x) = x^2 - x^3$	$[-1, 0]$
11. $f(x) = x(1 - x)^2$	$[0, 1]$
12. $f(x) = x^2(3 - x)$	$[0, 3]$

$\oplus$  In Exercises 13–16, use a program similar to that on page 366 to approximate the area of the region. How large must  $n$  be to obtain an approximation that is correct to within 0.01?

13.  $\int_0^4 (2x^2 + 3) dx$


14.  $\int_0^4 (2x^3 + 3) dx$

15.  $\int_1^2 (2x^2 - x + 1) dx$


16.  $\int_1^2 (x^3 - 1) dx$

In Exercises 17–20, use the Midpoint Rule with  $n = 4$  to approximate the area of the region. Compare your result with the exact area obtained with a definite integral.


17.  $f(y) = \frac{1}{4}y$ ,  $[2, 4]$




18.  $f(y) = 2y$ ,  $[0, 2]$



19.  $f(y) = y^2 + 1$ ,  $[0, 4]$




20.  $f(y) = 4y - y^2$ ,  $[0, 4]$



**Trapezoidal Rule** In Exercises 21 and 22, use the Trapezoidal Rule with  $n = 8$  to approximate the definite integral. Compare the result with the exact value and the approximation obtained with  $n = 8$  and the Midpoint Rule. Which approximation technique appears to be better? Let  $f$  be continuous on  $[a, b]$  and let  $n$  be the number of equal subintervals (see figure). Then the Trapezoidal Rule for approximating  $\int_a^b f(x) dx$  is

$$\frac{b-a}{2n} [f(x_0) + 2f(x_1) + \cdots + 2f(x_{n-1}) + f(x_n)].$$



21.  $\int_0^2 x^3 dx$

22.  $\int_1^3 \frac{1}{x^2} dx$

In Exercises 23–26, use the Trapezoidal Rule with  $n = 4$  to approximate the definite integral.

23.  $\int_0^2 \frac{1}{x+1} dx$

24.  $\int_0^4 \sqrt{1+x^2} dx$

25.  $\int_{-1}^1 \frac{1}{x^2+1} dx$

26.  $\int_1^5 \frac{\sqrt{x-1}}{x} dx$

⊕ In Exercises 27 and 28, use a computer or programmable calculator to approximate the definite integral using the Midpoint Rule and the Trapezoidal Rule for  $n = 4, 8, 12, 16,$  and  $20$ .

27.  $\int_0^4 \sqrt{2+3x^2} dx$

28.  $\int_0^2 \frac{5}{x^3+1} dx$

In Exercises 29 and 30, use the Trapezoidal Rule with  $n = 10$  to approximate the area of the region bounded by the graphs of the equations.


29.  $y = \sqrt{\frac{x^3}{4-x}}$ ,  $y = 0$ ,  $x = 3$

30.  $y = x\sqrt{\frac{4-x}{4+x}}$ ,  $y = 0$ ,  $x = 4$

31. **Velocity and Acceleration** The table lists the velocity  $v$  (in feet per second) of an accelerating car over a 20-second interval. Use the Trapezoidal Rule to approximate the distance in feet that the car travels during the 20 seconds. (The distance is given by  $s = \int_0^{20} v dt$ .)

Time, $t$	0	5	10	15	20
Velocity, $v$	0.0	29.3	51.3	66.0	73.3

32. **Surface Area** To estimate the surface area of a pond, a surveyor takes several measurements, as shown in the figure. Estimate the surface area of the pond using (a) the Midpoint Rule and (b) the Trapezoidal Rule.



⊕ 33. **Numerical Approximation** Use the Midpoint Rule and the Trapezoidal Rule with  $n = 4$  to approximate  $\pi$  where

$$\pi = \int_0^1 \frac{4}{1+x^2} dx.$$

Then use a graphing utility to evaluate the definite integral. Compare all of your results.

## The Disk


Another volume of type of the begin with and piston

As s plane reg

To d a continu area of th Figure 5. disks, ea revolving volumes can see th the Disk

## The C

The v graph



Approxim: FIGURE

**PREREQUISITE  
REVIEW 5.7**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, solve for  $x$ .

1.  $x^2 = 2x$

3.  $x = -x^3 + 5x$

5.  $-x + 4 = \sqrt{4x - x^2}$

2.  $-x^2 + 4x = x^2$

4.  $x^2 + 1 = x + 3$

6.  $\sqrt{x-1} = \frac{1}{2}(x-1)$

In Exercises 7–10, evaluate the integral.

7.  $\int_0^2 2e^{2x} dx$

9.  $\int_0^2 x\sqrt{x^2+1} dx$

8.  $\int_{-1}^3 \frac{2x+1}{x^2+x+2} dx$



10.  $\int_1^5 \frac{(\ln x)^2}{x} dx$

**EXERCISES 5.7**

In Exercises 1–16, find the volume of the solid formed by revolving the region bounded by the graph(s) of the equation(s) about the  $x$ -axis.



1.  $y = \sqrt{4-x^2}$

2.  $y = x^2$



3.  $y = \sqrt{x}$

4.  $y = \sqrt{4-x^2}$



5.  $y = 4 - x^2$ ,  $y = 0$

6.  $y = x$ ,  $y = 0$ ,  $x = 4$

7.  $y = 1 - \frac{1}{4}x^2$ ,  $y = 0$

8.  $y = x^2 + 1$ ,  $y = 5$

9.  $y = -x + 1$ ,  $y = 0$ ,  $x = 0$

10.  $y = x$ ,  $y = e^{x-1}$ ,  $x = 0$

11.  $y = \sqrt{x} + 1$ ,  $y = 0$ ,  $x = 0$ ,  $x = 9$

12.  $y = \sqrt{x}$ ,  $y = 0$ ,  $x = 4$

13.  $y = 2x^2$ ,  $y = 0$ ,  $x = 2$

14.  $y = \frac{1}{x}$ ,  $y = 0$ ,  $x = 1$ ,  $x = 3$

15.  $y = e^x$ ,  $y = 0$ ,  $x = 0$ ,  $x = 1$

16.  $y = x^2$ ,  $y = 4x - x^2$

In Exercises 17–24, find the volume of the solid formed by revolving the region bounded by the graph(s) of the equation(s) about the  $y$ -axis.

17.  $y = x^2$ ,  $y = 4$ ,  $0 \leq x \leq 2$



18.  $y = \sqrt{16-x^2}$ ,  $y = 0$ ,  $0 \leq x \leq 4$

19.  $x = 1 - \frac{1}{2}y$ ,  $x = 0$ ,  $y = 0$

20.  $x = y(y-1)$ ,  $x = 0$

21.  $y = x^{2/3}$

22.  $x = -y^2 + 4y$



Distance (in meters)

32. Volume  
revoluti

$y =$   
about  $t$   
measur

23.  $y = \sqrt{4-x}$   
24.  $y = 4$ ,  $y =$

25. Volume  
revolved at  
volume of  $t$

26. Volume  
revolved at  
volume of  $t$

27. Volume  
of a right ci  
base and  $h$

28. Volume  
of a sphere

29. Volume  
 $9x^2 +$

is revolve  
(shaped li  
spheroid.

30. Volume  
 $9x^2 +$

is revolve  
(shaped li

31. Volume  
by revol  
 $y = \frac{1}{8}x^2$   
and  $y$  are  
of the tan



23.  $y = \sqrt{4-x}$ ,  $y = 0$ ,  $x = 0$

24.  $y = 4$ ,  $y = 0$ ,  $x = 2$ ,  $x = 0$

25. **Volume** The line segment from  $(0, 0)$  to  $(6, 3)$  is revolved about the  $x$ -axis to form a cone. What is the volume of the cone?

26. **Volume** The line segment from  $(0, 0)$  to  $(4, 2)$  is revolved about the  $y$ -axis to form a cone. What is the volume of the cone?

27. **Volume** Use the Disk Method to verify that the volume of a right circular cone is  $\frac{1}{3}\pi r^2 h$ , where  $r$  is the radius of the base and  $h$  is the height.

28. **Volume** Use the Disk Method to verify that the volume of a sphere of radius  $r$  is  $\frac{4}{3}\pi r^3$ .

29. **Volume** The right half of the ellipse

$$9x^2 + 25y^2 = 225$$


is revolved about the  $y$ -axis to form an oblate spheroid (shaped like an M&M candy). Find the volume of the spheroid.

30. **Volume** The upper half of the ellipse

$$9x^2 + 16y^2 = 144$$

is revolved about the  $x$ -axis to form a prolate spheroid (shaped like a football). Find the volume of the spheroid.

31. **Volume** A tank on the wing of a jet airplane is modeled by revolving the region bounded by the graph of  $y = \frac{1}{8}x^2\sqrt{2-x}$  and the  $x$ -axis about the  $x$ -axis, where  $x$  and  $y$  are measured in meters (see figure). Find the volume of the tank.



32. **Volume** A soup bowl can be modeled as a solid of revolution formed by revolving the graph of

$$y = \sqrt{\frac{x}{2}} + 1, \quad 0 \leq x \leq 4$$

about the  $x$ -axis. Use this model, where  $x$  and  $y$  are measured in inches, to find the volume of the soup bowl.

33. **Biology** A pond is to be stocked with a species of fish. The food supply in 500 cubic feet of pond water can adequately support one fish. The pond is nearly circular, is 20 feet deep at its center, and has a radius of 200 feet. The bottom of the pond can be modeled by

$$y = 20[(0.005x)^2 - 1].$$

(a) How much water is in the pond?


(b) How many fish can the pond support?

34. **Modeling a Body of Water** A pond is approximately circular, with a diameter of 400 feet (see figure). Starting at the center, the depth of the water is measured every 25 feet and recorded in the table.

$x$	0	25	50	75	100
Depth	20	19	19	17	15

$x$	125	150	175	200
Depth	14	10	6	0

- (a) Use a graphing utility to plot the depths and graph the model of the pond's depth,  $y = 20 - 0.00045x^2$ .
- (b) Use the model in part (a) to find the pond's volume.
- (c) Use the result of part (b) to approximate the number of gallons of water in the pond ( $1 \text{ ft}^3 \approx 7.48 \text{ gal}$ ).



In Exercises 35 and 36, use a program similar to the one on page 366 to approximate the volume of a solid generated by revolving the region bounded by the graphs of the equations about the  $x$ -axis.

35.  $y = \sqrt[3]{x+1}$ ,  $y = 0$ ,  $x = 0$ ,  $x = 7$

36.  $y = \frac{10}{x^2+1}$ ,  $y = 0$ ,  $x = 0$ ,  $x = 3$