

**PREREQUISITE
REVIEW 6.1**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–8, evaluate the indefinite integral.

1. $\int 5 \, dx$

3. $\int x^{3/2} \, dx$

5. $\int 2x(x^2 + 1)^3 \, dx$

7. $\int 6e^{6x} \, dx$

2. $\int \frac{1}{3} \, dx$

4. $\int x^{2/3} \, dx$

6. $\int 3x^2(x^3 - 1)^2 \, dx$

8. $\int \frac{2}{2x + 1} \, dx$

In Exercises 9–12, simplify the expression.

9. $2x(x - 1)^2 + x(x - 1)$

11. $3(x + 7)^{1/2} - 2x(x + 7)^{-1/2}$

10. $6x(x + 4)^3 - 3x^2(x + 4)^2$

12. $(x + 5)^{1/3} - 5(x + 5)^{-2/3}$

EXERCISES 6.1

In Exercises 1–38, find the indefinite integral.

1. $\int (x - 2)^4 \, dx$

3. $\int \frac{2}{(t - 9)^2} \, dt$

5. $\int \frac{2t - 1}{t^2 - t + 2} \, dt$

7. $\int \sqrt{1 + x} \, dx$

9. $\int \frac{12x + 2}{3x^2 + x} \, dx$

11. $\int \frac{1}{(5x + 1)^3} \, dx$

13. $\int \frac{1}{\sqrt{x + 1}} \, dx$

15. $\int \frac{e^{3x}}{1 - e^{3x}} \, dx$

17. $\int \frac{2x}{e^{3x^2}} \, dx$

19. $\int \frac{x^2}{x - 1} \, dx$

21. $\int x\sqrt{x^2 + 4} \, dx$

23. $\int e^{5x} \, dx$

2. $\int (x + 5)^{3/2} \, dx$

4. $\int \frac{4}{(1 - t)^3} \, dt$

6. $\int \frac{2y^3}{y^4 + 1} \, dy$

8. $\int (3 + x)^{5/2} \, dx$

10. $\int \frac{6x^2 + 2}{x^3 + x} \, dx$

12. $\int \frac{1}{(3x + 1)^2} \, dx$

14. $\int \frac{1}{\sqrt{5x + 1}} \, dx$

16. $\int \frac{4e^{2x}}{1 + e^{2x}} \, dx$

18. $\int \frac{e^{\sqrt{x+1}}}{\sqrt{x+1}} \, dx$

20. $\int \frac{2x}{x - 4} \, dx$

22. $\int \frac{t}{\sqrt{1 - t^2}} \, dt$

24. $\int te^{t^2+1} \, dt$

25. $\int \frac{e^{-x}}{e^{-x} + 2} \, dx$

27. $\int \frac{x}{(x + 1)^4} \, dx$

29. $\int \frac{x}{(3x - 1)^2} \, dx$

31. $\int \frac{1}{\sqrt{t - 1}} \, dt$

33. $\int \frac{2\sqrt{t + 1}}{t} \, dt$

35. $\int \frac{x}{\sqrt{2x + 1}} \, dx$

37. $\int t^2\sqrt{1 - t} \, dt$

26. $\int \frac{e^x}{1 + e^x} \, dx$

28. $\int \frac{x^2}{(x + 1)^3} \, dx$

30. $\int \frac{5x}{(x - 4)^3} \, dx$

32. $\int \frac{1}{\sqrt{x + 1}} \, dx$

34. $\int \frac{6x + \sqrt{x}}{x} \, dx$

36. $\int \frac{x^2}{\sqrt{x - 1}} \, dx$

38. $\int y^2\sqrt[3]{y + 1} \, dy$

In Exercises 39–46, evaluate the definite integral.

39. $\int_0^4 \sqrt{2x + 1} \, dx$

41. $\int_0^1 3xe^{x^2} \, dx$

43. $\int_0^4 \frac{x}{(x + 4)^2} \, dx$

45. $\int_0^{0.5} x(1 - x)^3 \, dx$

40. $\int_2^4 \sqrt{4x + 1} \, dx$

42. $\int_0^2 e^{-2x} \, dx$

44. $\int_0^1 x(x + 5)^4 \, dx$

46. $\int_0^{0.5} x^2(1 - x)^3 \, dx$

In Exercise graphs of t region and

47. $y = x$

48. $y = x$

49. $y = x$

50. $y = x$

51. $y = -$

52. $y = \frac{2}{\sqrt{}}$

53. $y = x$

54. $y = x$

55. $y = -$

In Exercise graphs of t

56. $y = -$

In Exercise graphs of t

57. $y = -$

58. $y = -$

59. $y = x$

60. $y = \sqrt{}$

61. $f(x) =$

62. $f(x) =$

63. $f(x) =$

64. $f(x) =$

65. $f(x) =$

66. $f(x) =$

67. $f(x) =$

68. $f(x) =$

69. $f(x) =$

70. $f(x) =$

71. $f(x) =$

72. $f(x) =$

73. $f(x) =$

74. $f(x) =$

75. $f(x) =$

76. $f(x) =$

77. $f(x) =$

78. $f(x) =$

79. $f(x) =$

80. $f(x) =$

81. $f(x) =$

82. $f(x) =$

83. $f(x) =$

84. $f(x) =$

85. $f(x) =$

86. $f(x) =$

87. $f(x) =$

88. $f(x) =$

In Exercises 47–54, find the area of the region bounded by the graphs of the equations. Then use a graphing utility to graph the region and verify your answer.

47. $y = x\sqrt{x-3}$, $y = 0$, $x = 7$

48. $y = x\sqrt{2x+1}$, $y = 0$, $x = 4$

49. $y = x^2\sqrt{1-x}$, $y = 0$, $x = -3$

50. $y = x^2\sqrt{x+2}$, $y = 0$, $x = 7$

51. $y = \frac{x^2-1}{\sqrt{2x-1}}$, $y = 0$, $x = 1$, $x = 5$

52. $y = \frac{2x-1}{\sqrt{x+3}}$, $y = 0$, $x = \frac{1}{2}$, $x = 6$



53. $y = x\sqrt[3]{x+1}$, $y = 0$, $x = 0$, $x = 7$

54. $y = x\sqrt[3]{x-2}$, $y = 0$, $x = 2$, $x = 10$

In Exercises 55–58, find the area of the region bounded by the graphs of the equations.


55. $y = -x\sqrt{x+2}$, $y = 0$

56. $y = x\sqrt[3]{1-x}$, $y = 0$




57. $y^2 = x^2(1-x^2)$

(Hint: Find the area of the region bounded by $y = x\sqrt{1-x^2}$ and $y = 0$. Then multiply by 4.)



58. $y = 1/(1+\sqrt{x})$, $y = 0$, $x = 0$, $x = 4$



In Exercises 59 and 60, find the volume of the solid generated by revolving the region bounded by the graph(s) of the equation(s) about the x -axis.

59. $y = x\sqrt{1-x^2}$

60. $y = \sqrt{x}(1-x)^2$, $y = 0$

In Exercises 61 and 62, find the average amount by which the function f exceeds the function g on the interval.

61. $f(x) = \frac{1}{x+1}$, $g(x) = \frac{x}{(x+1)^2}$, $[0, 1]$

62. $f(x) = x\sqrt{4x+1}$, $g(x) = 2\sqrt{x^3}$, $[0, 2]$

63. Probability The probability of recall in an experiment is modeled by

$$P(a \leq x \leq b) = \int_a^b \frac{15}{4}x\sqrt{1-x} \, dx$$

where x is the percent of recall (see figure).

(a) What is the probability of recalling between 40% and 80%?

(b) What is the median percent recall? That is, for what value of b is $P(0 \leq x \leq b) = 0.5$?




Figure for 63




Figure for 64

64. Probability The probability of finding between a and b percent iron in ore samples is modeled by

$$P(a \leq x \leq b) = \int_a^b \frac{1155}{32}x^3(1-x)^{3/2} \, dx$$

(see figure). Find the probabilities that a sample will contain between (a) 0% and 25% and (b) 50% and 100% iron.

65. Meteorology During a two-week period in March in a small town near Lake Erie, the measurable snowfall S (in inches) on the ground can be modeled by

$$S(t) = t\sqrt{14-t}, \quad 0 \leq t \leq 14$$

where t represents the day.

- (a) Use a graphing utility to graph the function.
- (b) Find the average amount of snow on the ground during the two-week period.
- (c) Find the total snowfall over the two-week period.

66. Revenue A company sells a seasonal product that generates a daily revenue R (in dollars per year) modeled by

$$R = 0.06t^2(365-t)^{1/2} + 1250, \quad 0 \leq t \leq 365$$

where t represents the day.

- (a) Find the average daily revenue over a period of 1 year.
- (b) Describe a product whose seasonal sales pattern resembles the model. Explain your reasoning.

In Exercises 67 and 68, use a program similar to the Midpoint Rule program on page 366 with $n = 10$ to approximate the area of the region bounded by the graph(s) of the equation(s).

67. $y = \sqrt[3]{x}\sqrt{4-x}$, $y = 0$

68. $y^2 = x^2(1-x^2)$

**PREREQUISITE
REVIEW 6.2**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, find $f'(x)$.

1. $f(x) = \ln(x + 1)$

3. $f(x) = e^{x^3}$

5. $f(x) = x^2 e^x$

2. $f(x) = \ln(x^2 - 1)$

4. $f(x) = e^{-x^2}$

6. $f(x) = x e^{-2x}$

In Exercises 7–10, find the area between the graphs of f and g .

7. $f(x) = -x^2 + 4$, $g(x) = x^2 - 4$

8. $f(x) = -x^2 + 2$, $g(x) = 1$

9. $f(x) = 4x$, $g(x) = x^2 - 5$

10. $f(x) = x^3 - 3x^2 + 2$, $g(x) = x - 1$

EXERCISES 6.2

In Exercises 1–6, use integration by parts to find the indefinite integral.

1. $\int x e^{3x} dx$

2. $\int x e^{-x} dx$

3. $\int x^2 e^{-x} dx$

4. $\int x^2 e^{2x} dx$

5. $\int \ln 2x dx$

6. $\int \ln x^2 dx$

In Exercises 7–28, find the indefinite integral. (Hint: Integration by parts is not required for all the integrals.)

7. $\int e^{4x} dx$

8. $\int e^{-2x} dx$

9. $\int x e^{4x} dx$

10. $\int x e^{-2x} dx$

11. $\int x e^{x^2} dx$

12. $\int x^2 e^{x^3} dx$

13. $\int x^2 e^x dx$

14. $\int \frac{x}{e^x} dx$

15. $\int t \ln(t + 1) dt$

16. $\int x^3 \ln x dx$

17. $\int \frac{e^{1/t}}{t^2} dt$

18. $\int \frac{1}{x(\ln x)^3} dx$

19. $\int x(\ln x)^2 dx$

20. $\int \ln 3x dx$

21. $\int \frac{(\ln x)^2}{x} dx$

22. $\int \frac{1}{x \ln x} dx$

23. $\int x \sqrt{x-1} dx$

24. $\int \frac{x}{\sqrt{x-1}} dx$

25. $\int x(x+1)^2 dx$

26. $\int \frac{x}{\sqrt{2+3x}} dx$

27. $\int \frac{x e^{2x}}{(2x+1)^2} dx$

28. $\int \frac{x^3 e^{x^2}}{(x^2+1)^2} dx$

In Exercises 29–34, evaluate the definite integral.

29. $\int_0^1 x^2 e^x dx$


30. $\int_0^2 \frac{x^2}{e^x} dx$

31. $\int_1^e x^5 \ln x dx$

32. $\int_1^e 2x \ln x dx$

33. $\int_{-1}^0 \ln(x+2) dx$

34. $\int_0^1 \ln(1+2x) dx$

 In Exercises 35–38, find the area of the region bounded by the graphs of the equations. Then use a graphing utility to graph the region and verify your answer.

35. $y = x^3 e^x$, $y = 0$, $x = 0$, $x = 2$

36. $y = (x^2 - 1)e^x$, $y = 0$, $x = -1$, $x = 1$

37. $y = x^2 \ln x$, $y = 0$, $x = 1$, $x = e$

38. $y = \frac{\ln x}{x^2}$, $y = 0$, $x = 1$, $x = e$

In Exercises 39–42, find the indefinite integral using each specified method. Then write a brief statement explaining which method you prefer.

$$39. \int 2x\sqrt{2x-3} \, dx$$

(a) By parts, letting $dv = \sqrt{2x-3} \, dx$

(b) By substitution, letting $u = \sqrt{2x-3}$

$$40. \int x\sqrt{4+x} \, dx$$

(a) By parts, letting $dv = \sqrt{4+x} \, dx$

(b) By substitution, letting $u = \sqrt{4+x}$

$$41. \int \frac{x}{\sqrt{4+5x}} \, dx$$

(a) By parts, letting $dv = \frac{1}{\sqrt{4+5x}} \, dx$

(b) By substitution, letting $u = \sqrt{4+5x}$

$$42. \int x\sqrt{4-x} \, dx$$

(a) By parts, letting $dv = \sqrt{4-x} \, dx$

(b) By substitution, letting $u = \sqrt{4-x}$

In Exercises 43 and 44, use integration by parts to verify the formula.

$$43. \int x^n \ln x \, dx = \frac{x^{n+1}}{(n+1)^2} [-1 + (n+1) \ln x] + C,$$

$n \neq -1$

$$44. \int x^n e^{ax} \, dx = \frac{x^n e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} \, dx$$

In Exercises 45–48, use the results of Exercises 43 and 44 to find the indefinite integral.

$$45. \int x^2 e^{5x} \, dx$$

$$46. \int x e^{-3x} \, dx$$

$$47. \int x^{-2} \ln x \, dx$$

$$48. \int x^{1/2} \ln x \, dx$$

In Exercises 49–52, find the area of the region bounded by the graphs of the given equations.

$$49. y = xe^{-x}, y = 0, x = 4$$

$$50. y = \frac{1}{6}xe^{-x/3}, y = 0, x = 0, x = 3$$

$$51. y = x \ln x, y = 0, x = e$$


$$52. y = x^{-3} \ln x, y = 0, x = e$$

53. Given the region bounded by the graphs of $y = 2 \ln x$, $y = 0$, and $x = e$ (see figure), find

(a) the area of the region.

(b) the volume of the solid generated by revolving the region about the x -axis.


Figure for 53



54. Given the region bounded by the graphs of $y = xe^x$, $y = 0$, $x = 0$, and $x = 1$ (see figure), find

(a) the area of the region.

(b) the volume of the solid generated by revolving the region about the x -axis.



⊕ In Exercises 55–58, use a symbolic integration utility to evaluate the integral.

$$55. \int_0^2 t^3 e^{-4t} \, dt$$

$$56. \int_1^4 \ln x(x^2 + 4) \, dx$$

$$57. \int_0^5 x^4(25 - x^2)^{3/2} \, dx$$

$$58. \int_1^e x^9 \ln x \, dx$$

59. **Demand** A manufacturing company forecasts that the demand x (in units per year) for its product over the next 10 years can be modeled by $x = 500(20 + te^{-0.1t})$ for $0 \leq t \leq 10$, where t is the time in years.

⊕ (a) Use a graphing utility to decide whether the company is forecasting an increase or a decrease in demand over the decade.

(b) According to the model, what is the total demand over the next 10 years?

(c) Find the average annual demand during the 10-year period.

60. **Capital Campaign** The board of trustees of a college is planning a five-year capital gifts campaign to raise money for the college. The goal is to have an annual gift income I that is modeled by $I = 2000(375 + 68te^{-0.2t})$ for $0 \leq t \leq 5$, where t is the time in years.

⊕ (a) Use a graphing utility to decide whether the company is forecasting an increase or a decrease in demand over the decade.

(b) Find the total demand over the next 10 years.

(c) Find the average annual demand during the 10-year period.

61. **Learning** to memory

$$M = 1$$

where t is the time in years.

(a) the rate of learning

(b) the total learning

62. **Revenue**

revenue R product cost C

$$R = 4$$

where t is the time in years.

(a) Find the total revenue over the next 10 years.

(b) Find the total cost over the next 10 years.

(c) Find the profit over the next 10 years.

Present Value

the income c annual inflation rate r

$$63. c = 5000$$

$$64. c = 450,$$

$$65. c = 150,$$

$$66. c = 30,000$$

$$67. c = 1000$$

$$68. c = 5000$$

69. **Present Value** of an annuity

$$c = 1$$

(a) Find the present value of the annuity.

(b) Assume that the interest rate is 6% per year. Find the present value of the annuity.

- (a) Use a graphing utility to decide whether the board of trustees expects the gift income to increase or decrease over the five-year period.
- (b) Find the expected total gift income over the five-year period.
- (c) Determine the average annual gift income over the five-year period. Compare the result with the income given when $t = 3$.

61. Learning Theory A model for the ability M of a child to memorize, measured on a scale from 0 to 10, is

$$M = 1 + 1.6t \ln t, \quad 0 < t \leq 4$$

where t is the child's age in years. Find the average value of this model between

- (a) the child's first and second birthdays.
- (b) the child's third and fourth birthdays.

62. Revenue A company sells a seasonal product. The revenue R (in dollars per year) generated by sales of the product can be modeled by

$$R = 410.5t^2e^{-t/30} + 25,000, \quad 0 \leq t \leq 365$$

where t is the time in days.

- (a) Find the average daily receipts during the first quarter, which is given by $0 \leq t \leq 90$.
- (b) Find the average daily receipts during the fourth quarter, which is given by $274 \leq t \leq 365$.
- (c) Find the total daily receipts during the year.

Present Value In Exercises 63–68, find the present value of the income c (measured in dollars) over t_1 years at the given annual inflation rate r .

63. $c = 5000$, $r = 5\%$, $t_1 = 4$ years
64. $c = 450$, $r = 4\%$, $t_1 = 10$ years
65. $c = 150,000 + 2500t$, $r = 4\%$, $t_1 = 10$ years
66. $c = 30,000 + 500t$, $r = 7\%$, $t_1 = 6$ years
67. $c = 1000 + 50e^{t/2}$, $r = 6\%$, $t_1 = 4$ years
68. $c = 5000 + 25te^{t/10}$, $r = 6\%$, $t_1 = 10$ years

69. Present Value A company expects its income c during the next 4 years to be modeled by

$$c = 150,000 + 75,000t.$$

- (a) Find the actual income for the business over the 4 years.
- (b) Assuming an annual inflation rate of 4%, what is the present value of this income?

70. Present Value A professional athlete signs a three-year contract in which the earnings can be modeled by

$$c = 300,000 + 125,000t.$$

- (a) Find the actual value of the athlete's contract.
- (b) Assuming an annual inflation rate of 5%, what is the present value of the contract?

Future Value In Exercises 71 and 72, find the future value of the income (in dollars) given by $f(t)$ over t_1 years at the annual interest rate of r . If the function f represents a continuous investment over a period of t_1 years at an annual interest rate of r (compounded continuously), then the future value of the investment is given by

$$\text{Future value} = e^{rt_1} \int_0^{t_1} f(t)e^{-rt} dt.$$

71. $f(t) = 3000$, $r = 8\%$, $t_1 = 10$ years

72. $f(t) = 3000e^{0.05t}$, $r = 10\%$, $t_1 = 5$ years

73. Finance: Future Value Use the equation from Exercises 71 and 72 to calculate the following. (Source: Adapted from *Garman/Togue, Personal Finance, Fifth Edition*)

- (a) The future value of \$1200 saved each year for 10 years earning 7% interest.
- (b) A person who wishes to invest \$1200 each year finds one investment choice that is expected to pay 9% interest per year and another, riskier choice that may pay 10% interest per year. What is the difference in return (future value) if the investment is made for 15 years?

74. Consumer Awareness In 2004, the total cost to attend Pennsylvania State University for 1 year was estimated to be \$19,843. If your grandparents had continuously invested in a college fund according to the model

$$f(t) = 400t$$

for 18 years, at an annual interest rate of 10%, would the fund have grown enough to allow you to cover 4 years of expenses at Pennsylvania State University? (Source: Pennsylvania State University)

75. Use a program similar to the Midpoint Rule program on page 366 with $n = 10$ to approximate

$$\int_1^4 \frac{4}{\sqrt{x} + \sqrt[3]{x}} dx.$$

76. Use a program similar to the Midpoint Rule program on page 366 with $n = 12$ to approximate the volume of the solid generated by revolving the region bounded by the graphs of

$$y = \frac{10}{\sqrt{x}e^x}, \quad y = 0, \quad x = 1, \quad \text{and} \quad x = 4$$

about the x -axis.

**PREREQUISITE
REVIEW 6.3**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–8, factor the expression.

1. $x^2 - 16$

2. $x^2 - 25$

3. $x^2 - x - 12$

4. $x^2 + x - 6$

5. $x^3 - x^2 - 2x$

6. $x^3 - 4x^2 + 4x$

7. $x^3 - 4x^2 + 5x - 2$

8. $x^3 - 5x^2 + 7x - 3$

In Exercises 9–14, rewrite the improper rational expression as the sum of a proper rational expression and a polynomial.

9. $\frac{x^2 - 2x + 1}{x - 2}$

10. $\frac{2x^2 - 4x + 1}{x - 1}$

11. $\frac{x^3 - 3x^2 + 2}{x - 2}$

12. $\frac{x^3 + 2x - 1}{x + 1}$

13. $\frac{x^3 + 4x^2 + 5x + 2}{x^2 - 1}$

14. $\frac{x^3 + 3x^2 - 4}{x^2 - 1}$

EXERCISES 6.3

In Exercises 1–12, write the partial fraction decomposition for the expression.

1. $\frac{2(x + 20)}{x^2 - 25}$

2. $\frac{3x + 11}{x^2 - 2x - 3}$

3. $\frac{8x + 3}{x^2 - 3x}$

4. $\frac{10x + 3}{x^2 + x}$

5. $\frac{4x - 13}{x^2 - 3x - 10}$

6. $\frac{7x + 5}{6(2x^2 + 3x + 1)}$

7. $\frac{3x^2 - 2x - 5}{x^3 + x^2}$

8. $\frac{3x^2 - x + 1}{x(x + 1)^2}$

9. $\frac{x + 1}{3(x - 2)^2}$

10. $\frac{3x - 4}{(x - 5)^2}$

11. $\frac{8x^2 + 15x + 9}{(x + 1)^3}$

12. $\frac{6x^2 - 5x}{(x + 2)^3}$

19. $\int \frac{1}{2x^2 + x} dx$

20. $\int \frac{5}{x^2 + x - 6} dx$

21. $\int \frac{3}{x^2 + x - 2} dx$

22. $\int \frac{1}{4x^2 - 9} dx$

23. $\int \frac{5 - x}{2x^2 + x - 1} dx$

24. $\int \frac{x + 1}{x^2 + 4x + 3} dx$

25. $\int \frac{x^2 + 12x + 12}{x^3 - 4x} dx$

26. $\int \frac{3x^2 - 7x - 2}{x^3 - x} dx$

27. $\int \frac{x + 2}{x^2 - 4x} dx$

28. $\int \frac{4x^2 + 2x - 1}{x^3 + x^2} dx$

29. $\int \frac{4 - 3x}{(x - 1)^2} dx$

30. $\int \frac{x^4}{(x - 1)^3} dx$

31. $\int \frac{3x^2 + 3x + 1}{x(x^2 + 2x + 1)} dx$

32. $\int \frac{3x}{x^2 - 6x + 9} dx$

In Exercises 13–32, find the indefinite integral.

13. $\int \frac{1}{x^2 - 1} dx$

14. $\int \frac{9}{x^2 - 9} dx$

15. $\int \frac{-2}{x^2 - 16} dx$

16. $\int \frac{-4}{x^2 - 4} dx$

17. $\int \frac{1}{3x^2 - x} dx$

18. $\int \frac{3}{x^2 - 3x} dx$

In Exercises 33–40, evaluate the definite integral.

33. $\int_4^5 \frac{1}{9 - x^2} dx$

34. $\int_0^1 \frac{3}{2x^2 + 5x + 2} dx$

35. $\int_1^5 \frac{x - 1}{x^2(x + 1)} dx$

36. $\int_0^1 \frac{x^2 - x}{x^2 + x + 1} dx$

37. $\int_0^1 \frac{x^3}{x^2 - 2} dx$


38. $\int_0^1 \frac{x^3 - 1}{x^2 - 4} dx$

39. $\int_1^2 \frac{x^3 - 4x^2 - 3x + 3}{x^2 - 3x} dx$


40. $\int_2^4 \frac{x^4 - 4}{x^2 - 1} dx$

In Exercises 41–44, find the area of the shaded region.


41. $y = \frac{14}{16 - x^2}$




42. $y = \frac{-4}{x^2 - x - 6}$



43. $y = \frac{x + 1}{x^2 - x}$



44. $y = \frac{x^2 + 2x - 1}{x^2 - 4}$



In Exercises 45–48, write the partial fraction decomposition for the rational expression. Check your result algebraically. Then assign a value to the constant a and use a graphing utility to check the result graphically.

45. $\frac{1}{a^2 - x^2}$

46. $\frac{1}{x(x + a)}$

47. $\frac{1}{x(a - x)}$

48. $\frac{1}{(x + 1)(a - x)}$

In Exercises 49–52, use a graphing utility to graph the function. Then find the volume of the solid generated by revolving the region bounded by the graphs of the given equations about the x -axis by using the integration capabilities of a graphing utility and by integrating by hand using partial fraction decomposition.

49. $y = \frac{10}{x(x + 10)}$, $y = 0$, $x = 1$, $x = 5$

50. $y = \frac{-4}{(x + 1)(x - 4)}$, $y = 0$, $x = 0$, $x = 3$

51. $y = \frac{2}{x^2 - 4}$, $x = 1$, $x = -1$, $y = 0$

52. $y = \frac{25x}{x^2 + x - 6}$, $x = -2$, $x = 0$, $y = 0$

53. Biology A conservation organization releases 100 animals of an endangered species into a game preserve. The organization believes that the preserve has a capacity of 1000 animals and that the herd will grow according to a logistic growth model. That is, the size y of the herd will follow the equation

$$\int \frac{1}{y(1000 - y)} dy = \int k dt$$

where t is measured in years. Find this logistic curve. (To solve for the constant of integration C and the proportionality constant k , assume $y = 100$ when $t = 0$ and $y = 134$ when $t = 2$.) Use a graphing utility to graph your solution.

54. Health: Epidemic A single infected individual enters a community of 500 individuals susceptible to the disease. The disease spreads at a rate proportional to the product of the total number infected and the number of susceptible individuals not yet infected. A model for the time it takes for the disease to spread to x individuals is

$$t = 5010 \int \frac{1}{(x + 1)(500 - x)} dx$$

where t is the time in hours.

- (a) Find the time it takes for 75% of the population to become infected (when $t = 0$, $x = 1$).
- (b) Find the number of people infected after 100 hours.

55. Marketing After test-marketing a new menu item, a fast-food restaurant predicts that sales of the new item will grow according to the model

$$\frac{dS}{dt} = \frac{2t}{(t + 4)^2}$$

where t is the time in weeks and S is the sales (in thousands of dollars). Find the sales of the menu item at 10 weeks.

56. Biology One gram of a bacterial culture is present at time $t = 0$, and 10 grams is the upper limit of the culture's weight. The time required for the culture to grow to y grams is modeled by

$$kt = \int \frac{1}{y(10 - y)} dy$$

where y is the weight of the culture (in grams) and t is the time in hours.

(a) Verify th
by

$y =$

Use the

(b) Use the

Weight (in grams)
10
9
8
7
6
5
4
3
2
1

57. Revenue
for Symant
modeled by

$$R = \frac{41}{t}$$

where $t =$
from 1995
during this

58. Medicine
semester b
history of :

$$\frac{dN}{dt} = -$$

where N is

(a) Find th
with th
return

(b) If not
will th
tion of

59. Biology
mals of ar
organizati
increase at

$$\frac{dN}{dt} = -$$

where N is

(a) Use th
ulator


(b) Find th
es with

- (a) Verify that the weight of the culture at time t is modeled by


$$y = \frac{10}{1 + 9e^{-10kt}}$$

Use the fact that $y = 1$ when $t = 0$.

- (b) Use the graph to determine the constant k .



60. **Biology: Population Growth** The graph shows the logistic growth curves for two species of the single-celled *Paramecium* in a laboratory culture. During which time intervals is the rate of growth of each species increasing? During which time intervals is the rate of growth of each species decreasing? Which species has a higher limiting population under these conditions? (Source: Adapted from Levine/Miller, *Biology: Discovering Life, Second Edition*)



57. **Revenue** The revenue R (in millions of dollars per year) for Symantec Corporation from 1995 through 2003 can be modeled by

$$R = \frac{410t^2 + 28,490t + 28,080}{-6t^2 + 94t + 100}$$

where $t = 5$ corresponds to 1995. Find the total revenue from 1995 through 2003. Then find the average revenue during this time period. (Source: Symantec Corporation)

58. **Medicine** On a college campus, 50 students return from semester break with a contagious flu virus. The virus has a history of spreading at a rate of

$$\frac{dN}{dt} = \frac{100e^{-0.1t}}{(1 + 4e^{-0.1t})^2}$$

where N is the number of students infected after t days.

- (a) Find the model giving the number of students infected with the virus in terms of the number of days since returning from semester break.
- (b) If nothing is done to stop the virus from spreading, will the virus spread to infect half the student population of 1000 students? Explain your answer.
59. **Biology** A conservation organization releases 100 animals of an endangered species into a game preserve. The organization believes the population of the species will increase at a rate of

$$\frac{dN}{dt} = \frac{125e^{-0.125t}}{(1 + 9e^{-0.125t})^2}$$

where N is the population and t is the time in months.

- (a) Use the fact that $N = 100$ when $t = 0$ to find the population after 2 years.
- (b) Find the limiting size of the population as time increases without bound.

BUSINESS CAPSULE



Courtesy of Susie Wang/Aqua Dessa

While a math communications major at the University of California at Berkeley, Susie Wang began researching the idea of selling natural skin-care products. She used \$10,000 to start her company, Aqua Dessa, and uses word-of-mouth as an advertising tactic. Aqua Dessa products are used and sold at spas and exclusive cosmetics counters throughout the United States.

61. **Research Project** Use your school's library, the Internet, or some other reference source to research the opportunity cost of attending graduate school for 2 years to receive a Masters of Business Administration (MBA) degree rather than working for 2 years with a bachelor's degree. Write a short paper describing these costs.

**PREREQUISITE
REVIEW 6.5**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, find the indicated derivative.

1. $f(x) = \frac{1}{x}, f''(x)$

2. $f(x) = \ln(2x + 1), f^{(4)}(x)$

3. $f(x) = 2 \ln x, f^{(4)}(x)$

4. $f(x) = x^3 - 2x^2 + 7x - 12, f''(x)$

5. $f(x) = e^{2x}, f^{(4)}(x)$

6. $f(x) = e^{x^2}, f''(x)$

In Exercises 7 and 8, find the absolute maximum of f on the interval.

7. $f(x) = -x^2 + 6x + 9, [0, 4]$

8. $f(x) = \frac{8}{x^3}, [1, 2]$

In Exercises 9 and 10, solve for n .


9. $\frac{1}{4n^2} < 0.001$

10. $\frac{1}{16n^4} < 0.0001$


EXERCISES 6.5

In Exercises 1–12, use the Trapezoidal Rule and Simpson's Rule to approximate the value of the definite integral for the indicated value of n . Compare these results with the exact value of the definite integral. Round your answers to four decimal places.


1. $\int_0^2 x^2 dx, n = 4$




2. $\int_0^1 \left(\frac{x^2}{2} + 1\right) dx, n = 4$



3. $\int_0^2 (x^4 + 1) dx, n = 4$



4. $\int_1^2 \frac{1}{x} dx, n = 4$



5. $\int_0^2 x^3 dx, n = 8$

6. $\int_1^3 (4 - x^2) dx, n = 4$

7. $\int_1^2 \frac{1}{x} dx, n = 8$

8. $\int_1^2 \frac{1}{x^2} dx, n = 4$

9. $\int_0^4 \sqrt{x} dx, n = 8$

10. $\int_0^2 \sqrt{1+x} dx, n = 4$

11. $\int_0^1 \frac{1}{1+x} dx, n = 4$

12. $\int_0^2 x\sqrt{x^2+1} dx, n = 4$

In Exercises 13–20, approximate the integral using (a) the Trapezoidal Rule and (b) Simpson's Rule. (Round your answers to three significant digits.)

Definite Integral	Subdivisions
13. $\int_0^1 \frac{1}{1+x^2} dx$	$n = 4$
14. $\int_0^2 \frac{1}{\sqrt{1+x^3}} dx$	$n = 4$
15. $\int_0^1 \sqrt{1-x^2} dx$	$n = 4$
16. $\int_0^1 \sqrt{1-x^3} dx$	$n = 8$
17. $\int_0^2 e^{-x^2} dx$	$n = 2$
18. $\int_0^2 e^{-x^3} dx$	$n = 4$
19. $\int_0^3 \frac{1}{2-2x+x^2} dx$	$n = 6$
20. $\int_0^3 \frac{x}{2+x+x^2} dx$	$n = 6$

⊕ Present Value In Exercises 21 and 22, use a program similar to the Simpson's Rule program on page 430 with $n = 8$ to approximate the present value of the income $c(t)$ over t_1 years at the given annual interest rate r . Then use the integration capabilities of a graphing utility to approximate the present value. Compare the results. (Present value is defined in Section 6.2.)

21. $c(t) = 6000 + 200\sqrt{t}$, $r = 7\%$, $t_1 = 4$

22. $c(t) = 200,000 + 15,000\sqrt[3]{t}$, $r = 10\%$, $t_1 = 8$

⊕ Marginal Analysis In Exercises 23 and 24, use a program similar to the Simpson's Rule program on page 430 with $n = 4$ to approximate the change in revenue from the marginal revenue function dR/dx . In each case, assume that the number of units sold x increases from 14 to 16.

23. $\frac{dR}{dx} = 5\sqrt{8000 - x^3}$

24. $\frac{dR}{dx} = 50\sqrt{x}\sqrt{20-x}$

⊕ Probability In Exercises 25–28, use a program similar to the Simpson's Rule program on page 430 with $n = 6$ to approximate the indicated normal probability. The standard normal probability density function is

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

if x is chosen at random from a population with this density, then the probability that x lies in the interval $[a, b]$ is

$$P(a \leq x \leq b) = \int_a^b f(x) dx.$$

25. $P(0 \leq x \leq 1)$


26. $P(0 \leq x \leq 2)$

27. $P(0 \leq x \leq 4)$

28. $P(0 \leq x \leq 1.5)$

⊕ Surveying In Exercises 29 and 30, use a program similar to the Simpson's Rule program on page 430 to estimate the number of square feet of land in the lot, where x and y are measured in feet, as shown in the figures. In each case, the land is bounded by a stream and two straight roads.


29.



x	0	100	200	300	400	500
y	125	125	120	112	90	90

x	600	700	800	900	1000
y	95	88	75	35	0

30.



x	0	10	20	30	40	50	60
y	75	81	84	76	67	68	69

x	70	80	90	100	110	120
y	72	68	56	42	23	0

In Exercises 31 and (b) Simps

31. $\int_0^2 x^4 dx$

33. $\int_0^1 e^{x^3} dx$

In Exercises 3 error in the 0.0001 using

35. $\int_0^1 x^4 dx$

37. $\int_1^3 e^{2x} dx$

In Exercises 3 page 430 to a

39. $\int_1^4 x\sqrt{x}$

41. $\int_2^5 10xe^x$

43. Prove th mate the demonst

$$\int_0^1 x^2$$

44. Use a pr page 430 ated by 1

$y =$ about the

In Exercises 4 required arc length of f be

$$\int_b^a \sqrt{1 + f'(x)^2}$$

45. Arc Len feet long

$y =$

Use a pr page 43 cable. C using th integrati

In Exercises 31–34, use the error formulas to find bounds for the error in approximating the integral using (a) the Trapezoidal Rule and (b) Simpson's Rule. (Let $n = 4$.)

$$31. \int_0^2 x^4 dx \qquad 32. \int_0^1 \frac{1}{x+1} dx$$

$$33. \int_0^1 e^{x^3} dx \qquad 34. \int_0^1 e^{-x^2} dx$$

In Exercises 35–38, use the error formulas to find n such that the error in the approximation of the definite integral is less than 0.0001 using (a) the Trapezoidal Rule and (b) Simpson's Rule.

$$35. \int_0^1 x^4 dx \qquad 36. \int_1^3 \frac{1}{x} dx$$

$$37. \int_1^3 e^{2x} dx \qquad 38. \int_3^5 \ln x dx$$

In Exercises 39–42, use the program for Simpson's Rule given on page 430 to approximate the integral. Use $n = 100$.

$$39. \int_1^4 x\sqrt{x+4} dx \qquad 40. \int_1^4 x^2\sqrt{x+4} dx$$

$$41. \int_2^5 10xe^{-x} dx \qquad 42. \int_2^5 10x^2e^{-x} dx$$

43. Prove that Simpson's Rule is exact when used to approximate the integral of a cubic polynomial function, and demonstrate the result for

$$\int_0^1 x^3 dx, \quad n = 2.$$

44. Use a program similar to the Simpson's Rule program on page 430 with $n = 4$ to find the volume of the solid generated by revolving the region bounded by the graphs of

$$y = x\sqrt[3]{x+4}, \quad y = 0, \quad \text{and} \quad x = 4$$

about the x -axis.

In Exercises 45 and 46, use the definite integral below to find the required arc length. If f has a continuous derivative, then the arc length of f between the points $(a, f(a))$ and $(b, f(b))$ is


$$\int_a^b \sqrt{1 + [f'(x)]^2} dx.$$

45. **Arc Length** The suspension cable on a bridge that is 400 feet long is in the shape of a parabola whose equation is

$$y = \frac{x^2}{800} \quad (\text{see figure}).$$

Use a program similar to the Simpson's Rule program on page 430 with $n = 12$ to approximate the length of the cable. Compare this result with the length obtained by using the table of integrals in Section 6.4 to perform the integration.


Figure for 45



46. **Arc Length** A fleeing hare leaves its burrow $(0, 0)$ and moves due north (up the y -axis). At the same time, a pursuing lynx leaves from 1 yard east of the burrow $(1, 0)$ and always moves toward the fleeing hare (see figure). If the lynx's speed is twice that of the hare's, the equation of the lynx's path is

$$y = \frac{1}{3}(x^{3/2} - 3x^{1/2} + 2).$$

Find the distance traveled by the lynx by integrating over the interval $[0, 1]$.



47. **Medicine** A body assimilates a 12-hour cold tablet at a rate modeled by

$$\frac{dC}{dt} = 8 - \ln(t^2 - 2t + 4), \quad 0 \leq t \leq 12$$

where dC/dt is measured in milligrams per hour and t is the time in hours. Find the total amount of the drug absorbed into the body during the 12 hours.

48. **Medicine** The concentration M (in grams per liter) of a 6-hour allergy medicine in a body is modeled by

$$M = 12 - 4 \ln(t^2 - 4t + 6), \quad 0 \leq t \leq 6$$

where t is the time in hours since the allergy medication was taken. Find the average level of concentration in the body over the six-hour period.

49. **Consumer Trends** The rate of change S in the number of subscribers to a newly introduced magazine is modeled by

$$\frac{dS}{dt} = 1000t^2e^{-t}, \quad 0 \leq t \leq 6$$

where t is the time in years. Find the total increase in the number of subscribers during the first 6 years.

**PREREQUISITE
REVIEW 6.6**

The following warm-up exercises involve skills that were covered in earlier sections. You will use these skills in the exercise set for this section.

In Exercises 1–6, find the limit.

1. $\lim_{x \rightarrow 2} (2x + 5)$

2. $\lim_{x \rightarrow 1} \left(\frac{1}{x} + 2x^2 \right)$

3. $\lim_{x \rightarrow -4} \frac{x + 4}{x^2 - 16}$

4. $\lim_{x \rightarrow 0} \frac{x^2 - 2x}{x^3 + 3x^2}$

5. $\lim_{x \rightarrow 1} \frac{1}{\sqrt{x} - 1}$

6. $\lim_{x \rightarrow -3} \frac{x^2 + 2x - 3}{x + 3}$

In Exercises 7–10, evaluate the expression (a) when $x = b$ and (b) when $x = 0$.

7. $\frac{4}{3}(2x - 1)^3$

8. $\frac{1}{x - 5} + \frac{3}{(x - 2)^2}$

9. $\ln(5 - 3x^2) - \ln(x + 1)$



10. $e^{3x^2} + e^{-3x^2}$

EXERCISES 6.6

In Exercises 1–14, determine whether or not the improper integral converges. If it does, evaluate the integral.

1. $\int_0^{\infty} e^{-x} dx$

2. $\int_{-\infty}^0 e^{2x} dx$



3. $\int_1^{\infty} \frac{1}{x^2} dx$

4. $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$

5. $\int_0^{\infty} e^{x/3} dx$

6. $\int_0^{\infty} \frac{5}{e^{2x}} dx$

7. $\int_3^{\infty} \frac{x}{\sqrt{x^2 - 16}} dx$

8. $\int_{1/2}^{\infty} \frac{1}{\sqrt{2x - 1}} dx$

9. $\int_{-\infty}^0 e^{-x} dx$

10. $\int_{-\infty}^{-1} \frac{1}{x^2} dx$

11. $\int_1^{\infty} \frac{e^{\sqrt{x}}}{\sqrt{x}} dx$

12. $\int_{-\infty}^0 \frac{x}{x^2 + 1} dx$



13. $\int_{-\infty}^{\infty} 2xe^{-3x^2} dx$

14. $\int_{-\infty}^{\infty} x^2 e^{-x^3} dx$

In Exercises 15–18, determine the divergence or convergence of the improper integral. Evaluate the integral if it converges.

15. $\int_0^4 \frac{1}{\sqrt{x}} dx$

16. $\int_3^4 \frac{1}{\sqrt{x-3}} dx$



In Exercises

19. $\int_0^1 \frac{1}{1-x} dx$

21. $\int_0^9 \sqrt{x} dx$

23. $\int_0^1 \frac{1}{x^2} dx$

25. $\int_0^2 \sqrt[3]{x} dx$

26. $\int_0^2 \sqrt{x} dx$

27. $\int_3^4 \sqrt{x} dx$

28. $\int_3^5 \frac{1}{x^2} dx$

In Exercises the graphs solid gener:

29. $y = \frac{1}{x^2}$

30. $y = e^{-x}$

In Exercises a and n to c

$$\lim_{x \rightarrow \infty} x^n$$

x	
$x^n e^{-ax}$	


31. $a = 1,$

32. $a = 2,$


33. $a = \frac{1}{2},$

34. $a = \frac{1}{2},$

17. $\int_0^2 \frac{1}{(x-1)^{2/3}} dx$



18. $\int_0^2 \frac{1}{(x-1)^2} dx$



In Exercises 19–28, evaluate the improper integral.

19. $\int_0^1 \frac{1}{1-x} dx$

20. $\int_0^{27} \frac{5}{\sqrt[3]{x}} dx$

21. $\int_0^9 \frac{1}{\sqrt{9-x}} dx$

22. $\int_0^2 \frac{x}{\sqrt{4-x^2}} dx$

23. $\int_0^1 \frac{1}{x^2} dx$

24. $\int_0^1 \frac{1}{x} dx$

25. $\int_0^2 \frac{1}{\sqrt[3]{x-1}} dx$

26. $\int_0^2 \frac{1}{(x-1)^{4/3}} dx$

27. $\int_3^4 \frac{1}{\sqrt{x^2-9}} dx$

28. $\int_3^5 \frac{1}{x^2\sqrt{x^2-9}} dx$

In Exercises 29 and 30, (a) find the area of the region bounded by the graphs of the given equations and (b) find the volume of the solid generated by revolving the region about the x -axis.

29. $y = \frac{1}{x^2}, y = 0, x \geq 1$

30. $y = e^{-x}, y = 0, x \geq 0$

In Exercises 31–34, complete the table for the specified values of a and n to demonstrate that

$$\lim_{x \rightarrow \infty} x^n e^{-ax} = 0, \quad a > 0, n > 0.$$

x	1	10	25	50
$x^n e^{-ax}$				

31. $a = 1, n = 1$

32. $a = 2, n = 4$

33. $a = \frac{1}{2}, n = 2$

34. $a = \frac{1}{2}, n = 5$

In Exercises 35–38, use the results of Exercises 31–34 to evaluate the improper integral.

35. $\int_0^{\infty} x^2 e^{-x} dx$

36. $\int_0^{\infty} (x-1)e^{-x} dx$

37. $\int_0^{\infty} x e^{-2x} dx$

38. $\int_0^{\infty} x e^{-x} dx$

39. Present Value A business is expected to yield a continuous flow of profit at the rate of \$500,000 per year. If money will earn interest at the nominal rate of 9% per year compounded continuously, what is the present value of the business (a) for 20 years and (b) forever? (Present value is defined in Section 6.2.)

40. Present Value Repeat Exercise 39 for a farm that is expected to produce a profit of \$75,000 per year. Assume that money will earn interest at the nominal rate of 8% compounded continuously. (Present value is defined in Section 6.2.)

Capitalized Cost In Exercises 41 and 42, find the capitalized cost C of an asset (a) for $n = 5$ years, (b) for $n = 10$ years, and (c) forever. The capitalized cost is given by

$$C = C_0 + \int_0^n c(t)e^{-rt} dt$$

where C_0 is the original investment, t is the time in years, r is the annual interest rate compounded continuously, and $c(t)$ is the annual cost of maintenance. [Hint: For part (c), see Exercises 31–34.]

41. $C_0 = \$650,000, c(t) = 25,000, r = 10\%$

42. $C_0 = \$650,000, c(t) = \$25,000(1 + 0.08t), r = 12\%$

43. Women's Height The mean height of American women between the ages of 25 and 34 is 64.5 inches, and the standard deviation is 2.4 inches. Find the probability that a 25- to 34-year-old woman chosen at random is

(a) between 5 and 6 feet tall.

(b) 5 feet 8 inches or taller.

(c) 6 feet or taller.

[Source: U.S. National Center for Health Statistics.]

44. Quality Control A company manufactures wooden yardsticks. The lengths of the yardsticks are normally distributed with a mean of 36 inches and a standard deviation of 0.2 inch. Find the probability that a yardstick is

(a) longer than 35.5 inches.

(b) longer than 35.9 inches.

vergence of
erges.



6 CHAPTER REVIEW EXERCISES

In Exercises 1–12, use a basic integration formula to find the indefinite integral.

- | | |
|------------------------------------|---|
| 1. $\int dt$ | 2. $\int (x^2 + 2x - 1) dx$ |
| 3. $\int (x + 5)^3 dx$ | 4. $\int \frac{2}{(x - 1)^2} dx$ |
| 5. $\int e^{10x} dx$ | 6. $\int 3xe^{-x^2} dx$ |
| 7. $\int \frac{1}{5x} dx$ | 8. $\int \frac{2x^3 - x}{x^4 - x^2 + 1} dx$ |
| 9. $\int x\sqrt{x^2 + 4} dx$ | 10. $\int \frac{1}{\sqrt{2x - 9}} dx$ |
| 11. $\int \frac{2e^x}{3 + e^x} dx$ | 12. $\int (x^2 - 1)e^{x^3 - 3x} dx$ |

In Exercises 13–20, use substitution to find the indefinite integral.

- | | |
|-----------------------------------|---|
| 13. $\int x(x - 2)^3 dx$ | 14. $\int x(1 - x)^2 dx$ |
| 15. $\int x\sqrt{x + 1} dx$ | 16. $\int x^2\sqrt{x + 1} dx$ |
| 17. $\int 2x\sqrt{x - 3} dx$ | 18. $\int \frac{\sqrt{x}}{1 + \sqrt{x}} dx$ |
| 19. $\int (x + 1)\sqrt{1 - x} dx$ | 20. $\int \frac{x}{x - 1} dx$ |

In Exercises 21–24, use substitution to evaluate the definite integral. Use a symbolic integration utility to verify your answer.

21. $\int_2^3 x\sqrt{x - 2} dx$
22. $\int_2^3 x^2\sqrt{x - 2} dx$
23. $\int_1^3 x^2(x - 1)^3 dx$
24. $\int_{-3}^0 x(x + 3)^4 dx$


25. **Probability** The probability of recall in an experiment is found to be

$$P(a \leq x \leq b) = \int_a^b \frac{105}{16} x^2 \sqrt{1 - x} dx$$

where x represents the percent of recall (see figure).

- (a) Find the probability that a randomly chosen individual will recall 80% of the material.

(b) What is the median percent recall? That is, for what value of b is it true that $P(0 \leq x \leq b) = 0.5$?




26. **Probability** The probability of locating between a and b percent of oil and gas deposits in a region is

$$P(a \leq x \leq b) = \int_a^b \frac{140}{27} x^2 (1 - x)^{1/3} dx$$

(see figure).

- (a) Find the probability that between 40% and 60% of the deposits will be found.
 (b) Find the probability that between 0% and 50% of the deposits will be found.



27. **Profit** The net profits P (in millions of dollars per year) for Home Depot from 1997 through 2003 can be modeled by

$$P = -925.08 + 154.753t \ln t, \quad 7 \leq t \leq 13$$

where t is the time in years, with $t = 7$ corresponding to 1997. (Source: The Home Depot, Inc.)

- (a) Find the average net profit for the years 1997 through 2003.
 (b) Find the total net profit for the years 1997 through 2003.

28. **Medicine** after inge

$$E(t) =$$

where the

(a) Use a val 0

(b) Over decre:

(c) At wl effecti

(d) Deter over t

(e) At wh tivene: interv:

In Exercises 29 integral.

29. $\int \frac{\ln x}{\sqrt{x}} dx$

30. $\int \sqrt{x} \ln x dx$

31. $\int (x - 1)$

32. $\int \ln\left(\frac{x}{x + 1}\right) dx$

In Exercises 33 the indefinite integral your answer.

33. $\int 2x^2 e^{2x} dx$

Present Value the income give given annual in

35. $c(t) = 10,000$

36. $c(t) = 20,000$

37. $c(t) = 12,000$

38. $c(t) = 10,000$

39. **Economic** of each sce

(a) \$1000 and 15%

(b) A lotte over 2C

(Source: A Edition)

28. **Medicine** The effectiveness of a pain-killing drug t hours after ingestion can be modeled by

$$E(t) = te^{-0.4t}$$

where the effectiveness E is measured as a percent.

- (a) Use a graphing utility to graph the model over the interval $0 \leq t \leq 24$.
- (b) Over what intervals is the effectiveness increasing and decreasing?
- (c) At what time does the medication reach maximum effectiveness?
- (d) Determine the average effectiveness of the medication over the 24-hour period.
- (e) At what time does the medication have the same effectiveness as the average effectiveness for the entire interval?

In Exercises 29–32, use integration by parts to find the indefinite integral.

29. $\int \frac{\ln x}{\sqrt{x}} dx$

30. $\int \sqrt{x} \ln x dx$

31. $\int (x - 1)e^x dx$

32. $\int \ln\left(\frac{x}{x+1}\right) dx$

In Exercises 33 and 34, use integration by parts repeatedly to find the indefinite integral. Use a symbolic integration utility to verify your answer.

33. $\int 2x^2e^{2x} dx$

34. $\int (\ln x)^3 dx$

Present Value In Exercises 35–38, find the present value of the income given by $c(t)$ (measured in dollars) over t_1 years at the given annual inflation rate r .

35. $c(t) = 10,000$, $r = 4\%$, $t_1 = 5$ years
36. $c(t) = 20,000 + 1500t$, $r = 6\%$, $t_1 = 10$ years
37. $c(t) = 12,000t$, $r = 5\%$, $t_1 = 10$ years
38. $c(t) = 10,000 + 100e^{t/2}$, $r = 5\%$, $t_1 = 5$ years

39. **Economics: Present Value** Calculate the present values of each scenario.

- (a) \$1000 per year for 5 years at interest rates of 5%, 10%, and 15%
- (b) A lottery ticket that pays \$100,000 per year after taxes over 20 years, assuming an inflation rate of 8%

(Source: Adapted from Boyer/Melvin, *Economics*, Third Edition)

40. **Finance: Present Value** You receive \$1000 at the end of each year for the next 3 years to help with college expenses. Assuming an annual interest rate of 6%, what is the present value of that stream of payments? (Source: Adapted from Garman/Torgue, *Personal Finance*, Fifth Edition)

41. **Finance: Present Value** Determine the amount a person planning for retirement would need to deposit today to be able to withdraw \$6000 each year for the next 10 years from an account earning 6% interest. (Source: Adapted from Garman/Torgue, *Personal Finance*, Fifth Edition)

42. **Finance: Present Value** A person invests \$50,000 earning 6% interest. If \$6000 is withdrawn each year, use present value to determine how many years it will take for the fund to run out. (Source: Adapted from Garman/Torgue, *Personal Finance*, Fifth Edition)

In Exercises 43–48, use partial fractions to find the indefinite integral.

43. $\int \frac{1}{x(x+5)} dx$

44. $\int \frac{4x-2}{3(x-1)^2} dx$

45. $\int \frac{4x-13}{x^2-3x-10} dx$

46. $\int \frac{4x^2-x-5}{x^2(x+5)} dx$

47. $\int \frac{x^2}{x^2+2x-15} dx$


48. $\int \frac{x^2+2x-12}{x(x+3)} dx$

49. **Sales** A new product initially sells 1250 units per week. After 24 weeks, the number of sales increases to 6500. The sales can be modeled by logistic growth with a limit of 10,000 units per week.

- (a) Find a logistic growth model for the number of units.
- (b) Use the model to complete the table.


Time, t	0	3	6	12	24
Sales, y					

- (c) Use the graph shown below to approximate the time t when sales will be 7500.



50. Biology A conservation society has introduced a population of 300 ring-necked pheasants into a new area. After 5 years, the population has increased to 966. The population can be modeled by logistic growth with a limit of 2700 pheasants.

- Find a logistic growth model for the population of ring-necked pheasants.
- How many pheasants were present after 4 years?
- How long will it take to establish a population of 1750 pheasants?



In Exercises 51–56, use the table of integrals in Section 6.4 to evaluate the integral.

- | | |
|---|---|
| 51. $\int \frac{\sqrt{x^2 + 25}}{x} dx$ | 52. $\int \frac{1}{x(4 + 3x)} dx$ |
| 53. $\int \frac{1}{x^2 - 4} dx$ | 54. $\int x(\ln x^2)^2 dx$ |
| 55. $\int_0^3 \frac{x}{\sqrt{1+x}} dx$ | 56. $\int_1^3 \frac{1}{x^2 \sqrt{16-x^2}} dx$ |

In Exercises 57–60, use a reduction formula from the table of integrals in Section 6.4 to find the indefinite integral.


- | | |
|------------------------------------|-------------------------------------|
| 57. $\int \frac{\sqrt{1+x}}{x} dx$ | 58. $\int \frac{1}{(x^2 - 9)^2} dx$ |
| 59. $\int (x - 5)^3 e^{x-5} dx$ | 60. $\int (\ln x)^4 dx$ |

In Exercises 61–64, complete the square and then use the table of integrals in Section 6.4 to find the indefinite integral.


- | | |
|---------------------------------------|--|
| 61. $\int \frac{1}{x^2 + 4x - 21} dx$ | 62. $\int \frac{1}{x^2 - 8x - 52} dx$ |
| 63. $\int \sqrt{x^2 - 10x} dx$ | 64. $\int \frac{x}{\sqrt{x^4 + 6x^2 + 10}} dx$ |

In Exercises 65–68, use the Trapezoidal Rule to approximate the definite integral.


65. $\int_1^3 \frac{1}{x^2} dx, n = 4$




66. $\int_0^2 (x^2 + 1) dx, n = 4$



67. $\int_1^2 \frac{1}{1 + \ln x} dx, n = 4$




68. $\int_0^2 \frac{1}{\sqrt{1+x^3}} dx, n = 8$




In Exercises 69–72, use Simpson's Rule to approximate the definite integral.


69. $\int_1^2 \frac{1}{x^3} dx, n = 4$




70. $\int_1^2 x^3 dx, n = 4$



71. $\int_0^1 \frac{x^{3/2}}{2-x^2} dx, n = 4$



72. $\int_0^1 e^{x^2} dx, n = 6$



In Exercises 73–74, use the Trapezoidal Rule to approximate the error in approximating the definite integral.

73. $\int_0^2 e^{2x} dx,$

74. $\int_0^2 e^{2x} dx,$

In Exercises 75–76, use Simpson's Rule to approximate the error in approximating the definite integral.

75. $\int_2^4 \frac{1}{x-1} dx$


76. $\int_2^4 \frac{1}{x-1} dx$

In Exercises 77–78, use Simpson's Rule to approximate the error in approximating the definite integral.


77. $\int_0^\infty 4xe^{-2x} dx$



79. $\int_{-\infty}^0 \frac{1}{3x^2} dx$



81. $\int_0^4 \frac{1}{\sqrt{4x}} dx$



In Exercises 73 and 74, use the error formula to find bounds for the error in approximating the integral using the Trapezoidal Rule.

73. $\int_0^2 e^{2x} dx, n = 4$

74. $\int_0^2 e^{2x} dx, n = 8$

In Exercises 75 and 76, use the error formula to find bounds for the error in approximating the integral using Simpson's Rule.



75. $\int_2^4 \frac{1}{x-1} dx, n = 4$

76. $\int_2^4 \frac{1}{x-1} dx, n = 8$

In Exercises 77–84, evaluate the improper integral.



77. $\int_0^\infty 4xe^{-2x^2} dx$

78. $\int_{-\infty}^0 \frac{3}{(1-3x)^{2/3}} dx$





79. $\int_{-\infty}^0 \frac{1}{3x^2} dx$

80. $\int_0^\infty 2x^2e^{-x^3} dx$





81. $\int_0^4 \frac{1}{\sqrt{4x}} dx$

82. $\int_1^2 \frac{x}{16(x-1)^2} dx$



83. $\int_2^3 \frac{1}{\sqrt{x-2}} dx$

84. $\int_0^2 \frac{x+2}{(x-1)^2} dx$



85. **Present Value** You are considering buying a franchise that yields a continuous income stream of \$50,000 per year. Find the present value of the franchise (a) for 15 years and (b) forever. Assume that money earns 6% interest per year, compounded continuously.

86. **Capitalized Cost** A company invests \$1.5 million in a new manufacturing plant that will cost \$50,000 per year in maintenance. Find the capitalized cost for (a) 20 years and (b) forever. Assume that money earns 6% interest, compounded continuously.

87. **SAT Scores** In 2001, the Scholastic Aptitude Test (SAT) math scores for college-bound seniors roughly followed a normal distribution

$$y = 0.0035e^{-(x-514)^2/25,538}, \quad 200 \leq x \leq 800$$

where x is the SAT score for mathematics. Find the probability that a senior chosen at random had an SAT score (a) between 500 and 650, (b) 650 or better, and (c) 750 or better. (Source: College Board)

