

## THE NUMERICAL SOLUTION OF THE BIHARMONIC EQUATION BY CONFORMAL MAPPING\*

RAYMOND H. CHAN<sup>†</sup>, THOMAS K. DELILLO<sup>‡</sup>, AND MARK A. HORN<sup>‡</sup>

**Abstract.** The solution to the biharmonic equation in a simply connected region  $\Omega$  in the plane is computed in terms of the Goursat functions. The boundary conditions are conformally transplanted to the disk with a numerical conformal map. A linear system is obtained for the Taylor coefficients of the Goursat functions. The coefficient matrix of the linear system can be put in the form  $I + K$ , where  $K$  is the discretization of a compact operator.  $K$  can be thought of as the composition of a block Hankel matrix with a diagonal matrix. The compactness leads to clustering of eigenvalues, and the Hankel structure yields a matrix-vector multiplication cost of  $O(N \log N)$ . Thus, if the conjugate gradient method is applied to the system, then superlinear convergence will be obtained. Numerical results are given to illustrate the spectrum clustering and superlinear convergence.

**Key words.** biharmonic equation, numerical conformal mapping, Hankel matrices

**AMS subject classifications.** 30C30, 31A30, 65E05

**PII.** S1064827595292710

**1. Introduction.** Boundary value problems for the biharmonic equation in two dimensions arise in the computation of the Airy stress function for plane stress problems [KK], [Mik], [Musk], and in steady Stokes flow of highly viscous fluids [MT, Chap. 22], [Poz]. Integral equations methods are a popular choice for the numerical solution of these equations [GGMa], [MG], [K, and references therein], [Poz]. The application of conformal mapping to this problem, though classical, is less well known [KK], [Musk]. Unlike the Laplace equation, the biharmonic equation is not preserved under conformal transplantation. However, a biharmonic function and its boundary values can be represented in terms of the analytic Goursat functions, and this representation can be transplanted with a conformal map to a computational region, such as a disk, an ellipse, or an annulus, where the boundary value problem can be solved more easily.

In this paper, we consider simply connected regions with analytic boundaries and use the unit disk as our computational region. In our examples, the conformal map  $f$  from the unit disk to the region is either known explicitly or approximated numerically. The boundary conditions for the biharmonic function are then transplanted by  $f$  to the disk, and a linear system for the Taylor coefficients of the Goursat functions in the disk is obtained and solved efficiently by conjugate-gradient-like methods. If the boundary of the target region is smooth enough (analytic in our examples), the continuous problem can be posed as a compact operator acting on some appropriate Banach space. This will lead to a clustering of the spectrum, and hence to superlinear convergence.

---

\*Received by the editors October 2, 1995; accepted for publication (in revised form) February 26, 1996.

<http://www.siam.org/journals/sisc/18-6/29271.html>

<sup>†</sup>Department of Mathematics, The Chinese University of Hong Kong, Shatin, Hong Kong (rchan@math.cuhk.hk). The research of this author was partially supported by HKRGC grant CUHK316/94E.

<sup>‡</sup>Department of Mathematics and Statistics, Wichita State University, Wichita, KS 67260-0033 (delillo@twsvum.uc.twsu.edu, mhorn@twsvum.uc.twsu.edu). The research of these authors was partially supported by U. S. Department of Energy grant DE-FG02-92ER25124 and National Science Foundation EPSCoR grant OSR-9255223.

We expect to be able to generalize this work to cases where the conformal map  $f$  is a Faber series map from an ellipse, or a cross-shaped or spoke-like region as in [DE] and [DEP]. If the target region has elongated sections, the conformal map from the disk may be severely ill-conditioned, and an ellipse or cross-shaped region may provide a better computational region. In [GGMa] the Sherman–Lauricella equation is solved for spoke-like regions which provide difficult regions for plane stress and plane strain problems. We anticipate that our Faber series methods may have advantages for such highly distorted regions. In cases for which the target region is not too distorted, so that the map from the disk is not too severely ill-conditioned, our method may also have some advantages. For instance, if several boundary value problems have to be solved for the same region, so that the conformal map only has to be computed once, our method, which is based on the fast Fourier transform (FFT), will give accurate answers in  $O(N \log N)$  for moderate-sized  $N$ . The methods in [GGMa] use the fast multipole method, which costs only  $O(N)$ , but with a large constant, so that large  $N$  are required in practice for it to be faster than the FFT. Below, we will use the FFT-based numerical conformal mapping method given in [Weg]. Introductions to numerical conformal mapping can be found in [Ga] and [He].

The outline of the paper is as follows. In section 2, we discuss the solution of boundary value problems for the biharmonic equation in terms of Goursat functions and the conformal map from the disk to the plane region. In section 3, we discuss the special structure of the exact linear system. We will see that the coefficient matrix of the (infinite) linear system is of the form  $I + HD$ , where  $I$  is the identity matrix,  $D$  is a diagonal matrix, and  $H$  is a block Hankel matrix. (A *Hankel matrix* is constant on the antidiagonals.) It will be seen that  $HD$  actually can be represented as a compact operator with a one-dimensional null space. This system can be symmetrized and solved (up to the null vector) using the conjugate gradient method. In section 4, we formulate the discrete problem. We will show how the conjugate gradient method is applied to the discrete system and how the matrix-vector multiplication can be carried out in  $O(N \log N)$ . In section 5, we give several numerical examples which illustrate the spectrum clustering, the superlinear convergence, and the discretization error.

**2. The biharmonic equation.** Here we will follow the presentation in [KK] and [Musk]. We wish to find the Airy stress function  $u$  for a simply connected region  $\Omega$  with a smooth boundary  $\Gamma$  in the  $\zeta$ -plane. Then  $u$  satisfies the biharmonic equation

$$\Delta^2 u = 0$$

for  $\zeta = \eta + i\mu \in \Omega$ . The two fundamental boundary value problems in elasticity seek to find  $u$ , given the external stresses or external displacements on the boundary  $\Gamma$ . Both of these problems amount to specifying

$$u_\eta = G_1 \quad \text{and} \quad u_\mu = G_2$$

on  $\Gamma$ . The function  $u$  can be represented as

$$u(\zeta) = \operatorname{Re}(\bar{\zeta}\phi(\zeta) + \chi(\zeta)),$$

where  $\phi(\zeta)$  and  $\chi(\zeta)$  are analytic functions in  $\Omega$  known as the *Goursat functions*. Letting  $G = G_1 + iG_2$ , the boundary conditions for the first fundamental problem become

$$(1) \quad \phi(\zeta) + \zeta\overline{\phi'(\zeta)} + \overline{\psi(\zeta)} = G(\zeta), \quad \zeta \in \Gamma,$$

where  $\psi(\zeta) = \chi'(\zeta)$ . The second fundamental problem leads to similar conditions. For simplicity, in this paper, we will only concentrate on the first boundary conditions (1).

We remark that  $\phi(\zeta)$  and  $\psi(\zeta)$  are not unique. In fact, if  $\phi(\zeta)$  and  $\psi(\zeta)$  represent any solution of the problem, then so does  $\phi(\zeta) + Ci\zeta + \gamma$  and  $\psi(\zeta) + \gamma'$ , where  $C \in \mathbf{R}$ ,  $\gamma \in \mathbf{C}$ , and  $\gamma' \in \mathbf{C}$ . Thus, the constants  $C$  and  $\gamma$  must be specified for uniqueness of  $\phi$ . These constants are determined below.

The problem at this point is to find  $\phi$  and  $\psi$  analytic in  $\Omega$  and satisfying (1). One approach is to represent  $\phi$  and  $\psi$  as Cauchy-type integrals of a density function on  $\Gamma$ . This leads to the *Sherman-Lauricella equation*, a Fredholm integral equation for the density function which can be solved efficiently by the fast multipole method [GGMa]. In this paper, we propose to solve it by using numerical conformal mapping coupled with the conjugate gradient method.

Let  $\zeta = f(z)$  be the conformal map from the unit disk to  $\Omega$ , fixing  $f(0) = 0 \in \Omega$ . Then with  $d(z) := f(z)/\overline{f'(z)}$ ,  $\phi(z) := \phi(f(z))$ ,  $\psi(z) := \psi(f(z))$ , and  $G(z) := G(f(z))$ , equation (1) transplants to the disk as

$$(2) \quad \phi(z) + d(z)\overline{\phi'(z)} + \overline{\psi(z)} = G(z), \quad |z| = 1.$$

Let

$$\phi(z) = \sum_{k=1}^{\infty} a_k z^k \quad \text{and} \quad \psi(z) = \sum_{k=0}^{\infty} b_k z^k.$$

Notice that the sum for  $\phi$  begins at  $k = 1$ . This fixes the constant  $\gamma$  mentioned above for uniqueness by requiring  $\phi(0) = a_0 = 0$ . After transplanting to the disk, the other constant is determined by setting  $\text{Im}(a_1/f'(0)) = 0$ .

The problem is to find the  $a_k$ 's and the  $b_k$ 's. For  $|z| = 1$ , define the Fourier series

$$d(z) := f(z)/\overline{f'(z)} = \sum_{k=-\infty}^{\infty} h_k z^k, \quad G(z) = \sum_{k=-\infty}^{\infty} A_k z^k.$$

Substituting into (2) gives a linear system of equations for the  $a_k$ 's and  $b_k$ 's,

$$(3) \quad a_j + \sum_{k=1}^{\infty} k \bar{a}_k h_{k+j-1} = A_j, \quad j = 1, 2, 3, \dots,$$

$$(4) \quad \bar{b}_j + \sum_{k=1}^{\infty} k \bar{a}_k h_{k-j-1} = A_{-j}, \quad j = 0, 1, 2, \dots$$

If (3) is solved for the  $a_k$ 's, then the  $b_k$ 's can be easily computed from (4). Thus, in this paper, we will concentrate on an efficient method for solving (3).

There is also a moment condition to be satisfied by the data. After transplantation to the disk, this condition can be stated as  $\text{Re}[\int_{|z|=1} G(z)\overline{f'(z)}dz] = 0$ . This moment condition will assure the existence of a solution. Our assumption is that all data studied in this paper satisfy this equation.

Before proceeding, it should be noted that if our boundary data corresponds to  $G = 0$ , then the only possible (nonzero) choice for  $\phi$  is  $\phi(z) = Cif(z)$ , for some nonzero  $C \in \mathbf{R}$ . This implies that the null space corresponding to the infinite system in (3) is one-dimensional, and the eigenvector spanning this space is given by  $a_k = ic_k, k = 1, 2, 3, \dots$ , where  $f(z) = \sum_{k=1}^{\infty} c_k z^k$ .

**3. Compact operators.** Taking real and imaginary parts of equation (3) gives us

$$(5) \quad \alpha_j + \sum_{k=1}^{\infty} k(\eta_{k+j-1}\alpha_k + \gamma_{k+j-1}\beta_k) = B_j, \quad j = 1, 2, 3, \dots,$$

$$(6) \quad \beta_j + \sum_{k=1}^{\infty} k(\gamma_{k+j-1}\alpha_k - \eta_{k+j-1}\beta_k) = C_j, \quad j = 1, 2, 3, \dots,$$

where we have used the notation  $a_k = \alpha_k + i\beta_k$ ,  $h_k = \eta_k + i\gamma_k$ , and  $A_k = B_k + iC_k$ . For visualization purposes, we combine equations (5) and (6) into a doubly infinite matrix equation in which the two sums are combined into a block Hankel matrix composed with a diagonal matrix. In fact, (5) and (6) can be written as

$$(7) \quad (I_{\infty} + H_{r,\infty}D_{\infty})\underline{\alpha} + H_{i,\infty}D_{\infty}\underline{\beta} = \underline{B},$$

$$(8) \quad (I_{\infty} - H_{r,\infty}D_{\infty})\underline{\beta} + H_{i,\infty}D_{\infty}\underline{\alpha} = \underline{C},$$

so that

$$(9) \quad \left( \begin{pmatrix} I_{\infty} & 0 \\ 0 & I_{\infty} \end{pmatrix} + \begin{pmatrix} H_{r,\infty} & H_{i,\infty} \\ H_{i,\infty} & -H_{r,\infty} \end{pmatrix} \begin{pmatrix} D_{\infty} & 0 \\ 0 & D_{\infty} \end{pmatrix} \right) \begin{pmatrix} \underline{\alpha} \\ \underline{\beta} \end{pmatrix} = \begin{pmatrix} \underline{B} \\ \underline{C} \end{pmatrix},$$

where  $\underline{\alpha} = (\alpha_1, \alpha_2, \dots)^T$ ,  $\underline{\beta} = (\beta_1, \beta_2, \dots)^T$ ,  $\underline{B} = (B_1, B_2, \dots)^T$ ,  $\underline{C} = (C_1, C_2, \dots)^T$ ,  $I_{\infty}$  is the infinite identity matrix,  $D_{\infty} = \text{diag}(1, 2, \dots)$ ,  $H_{r,\infty}$  is an infinite Hankel matrix generated by the  $\eta_k$ , and  $H_{i,\infty}$  is an infinite Hankel matrix generated by the  $\gamma_k$ .

Now suppose  $(\underline{\alpha}, \underline{\beta})$  represents a solution to (5), (6). Define

$$\underline{x} = \begin{pmatrix} D_{\infty}^{1/2}\underline{\alpha} \\ D_{\infty}^{1/2}\underline{\beta} \end{pmatrix}, \quad \underline{r} = \begin{pmatrix} D_{\infty}^{1/2}\underline{B} \\ D_{\infty}^{1/2}\underline{C} \end{pmatrix}.$$

Then (9) can be written as

$$(10) \quad (I_{\infty} + M_{\infty})\underline{x} = \underline{r},$$

where  $M_{\infty}$  is given by

$$M_{\infty} = \begin{pmatrix} M_{r,\infty} & M_{i,\infty} \\ M_{i,\infty} & -M_{r,\infty} \end{pmatrix} = \begin{pmatrix} D_{\infty}^{1/2}H_{r,\infty}D_{\infty}^{1/2} & D_{\infty}^{1/2}H_{i,\infty}D_{\infty}^{1/2} \\ D_{\infty}^{1/2}H_{i,\infty}D_{\infty}^{1/2} & -D_{\infty}^{1/2}H_{r,\infty}D_{\infty}^{1/2} \end{pmatrix}.$$

Note that  $M_{\infty}$  is symmetric. We would now like to justify the formal manipulations above and show that  $M_{\infty}$  is a compact operator. This will require the following two preliminary lemmas.

LEMMA 1. *Let  $f$  be a conformal map from the unit disk to the region  $\Omega$  with boundary  $\Gamma$ . Let  $\Gamma$  be analytic and*

$$f(e^{i\theta})/\overline{f'(e^{i\theta})} = \sum_{k=-\infty}^{\infty} h_k e^{ik\theta}.$$

*Then there exists a  $C > 0$  and an  $R < 1$  such that*

$$|h_k| \leq CR^{|k|}.$$

*Proof.* Since  $\Gamma$  is analytic,  $f$  extends as a bounded, analytic function with  $f'(z) \neq 0$  for  $|z| \leq 1/R$  for some  $R < 1$ . Let

$$f(z) = \sum_{k=1}^{\infty} c_k z^k \quad \text{and} \quad 1/f'(z) = \sum_{j=0}^{\infty} d_j z^j.$$

Then there is a  $c$  such that  $|c_k|, |d_k| \leq cR^k$ . Further, we have that

$$\begin{aligned} f(e^{i\theta})/\overline{f'(e^{i\theta})} &= \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} c_k \bar{d}_j e^{i(k-j)\theta} \\ &= \sum_{l=1}^{\infty} \sum_{j=0}^{\infty} c_{l+j} \bar{d}_j e^{il\theta} + \sum_{l=0}^{\infty} \sum_{j=l+1}^{\infty} c_{j-l} \bar{d}_j e^{-il\theta} \\ &= \sum_{l=1}^{\infty} \sum_{j=0}^{\infty} c_{l+j} \bar{d}_j e^{il\theta} + \sum_{l=0}^{\infty} \sum_{j=1}^{\infty} c_j \bar{d}_{l+j} e^{-il\theta}. \end{aligned}$$

And so,

$$|h_l| = \left| \sum_{j=0}^{\infty} c_{l+j} \bar{d}_j \right| \leq \sum_{j=0}^{\infty} |c_{l+j}| |\bar{d}_j| \leq cR^l \sum_{j=0}^{\infty} R^{2j} = \frac{cR^l}{1-R^2} = CR^l, \quad l \geq 1.$$

Similarly,

$$|h_{-l}| = \left| \sum_{j=1}^{\infty} c_j \bar{d}_{l+j} \right| \leq CR^l, \quad l \geq 0. \quad \square$$

Next we show that the entries of  $M_{r,\infty}$  and  $M_{i,\infty}$  also decay exponentially fast.

LEMMA 2. *Under the assumptions of Lemma 1, the  $(j, k)$ th entries of  $M_{r,\infty}$  and  $M_{i,\infty}$  decay like  $cr^{|j+k|}$  for some  $c > 0$  and  $r < 1$ .*

*Proof.* We will prove the case for  $M_{r,\infty}$ . The case for  $M_{i,\infty}$  follows similarly. Let  $m_{k,j}$  denote the  $(k, j)$ th entry of  $M_{r,\infty}$ . Then we must have  $m_{k,j} = \sqrt{kj}\eta_{k+j-1}$ . Therefore

$$(11) \quad |m_{k,j}| = \sqrt{kj}|h_{k+j-1}| \leq C\sqrt{kj}R^{|k+j|}.$$

Let  $r = (1 + R)/2 < 1$ . Since

$$\lim_{x \rightarrow \infty} \frac{1}{\sqrt{x}} \left(\frac{r}{R}\right)^x = \infty,$$

there exists an  $l_0 \geq 0$  such that

$$\sqrt{l}R^l \leq r^l \quad \forall l \geq l_0.$$

Let

$$c = \max_{0 \leq l \leq l_0} \left\{ \sqrt{l} \left(\frac{R}{r}\right)^l \right\};$$

we then see that

$$\sqrt{l}R^l \leq cr^l \quad \forall l \geq 0.$$

The lemma now follows directly from (11). □

Lemma 2 gives us the following theorem and corollary.

**THEOREM 1.**  $M_{r,\infty} : l^1 \rightarrow l^1$  and  $M_{i,\infty} : l^1 \rightarrow l^1$  are compact operators, where for  $\underline{y} \in l^1$ ,

$$M_{r,\infty}\underline{y} = \sum_{k=1}^{\infty} \sqrt{kj}\eta_{k+j-1}y_k, \quad j = 1, 2, \dots,$$

$$M_{i,\infty}\underline{y} = \sum_{k=1}^{\infty} \sqrt{kj}\gamma_{k+j-1}y_k, \quad j = 1, 2, \dots$$

*Proof.* We will prove the theorem for  $M_{r,\infty}$ . As above,  $M_{i,\infty}$  follows similarly. Define the finite rank operators  $\{M_{r,n}\} = \{D_n^{1/2}H_{r,n}D_n^{1/2}\}$  by

$$M_{r,n}\underline{y} = \sum_{k=1}^n \sqrt{kj}\eta_{k+j-1}y_k, \quad j = 1, 2, \dots, n,$$

for all  $\underline{y} = (y_1, y_2, \dots) \in l^1$ . The goal is to show that  $M_{r,\infty}$  can be approximated uniformly by these finite rank operators. (Then, e.g., a version of Theorem 4.4c [Con, p. 41] for Banach spaces shows that  $M_{r,\infty}$  is itself compact.) If  $A = (a_{kj})$  is an infinite matrix, then the induced  $l^1$  operator norm is given by

$$\|A\|_{l^1} = \sup_j \sum_{k=1}^{\infty} |a_{kj}|;$$

see, e.g., [Con, p. 171, prob. 8]. From the geometric decay of Lemma 2 we may write

$$\sum_{k=1}^{\infty} |m_{k,j}| \leq Cr^j \sum_{k=1}^{\infty} r^k \leq C_1r^j, \quad j \geq 1.$$

Consequently,

$$\begin{aligned} \|M_{r,\infty} - M_{r,n}\|_{l^1} &= \sup \left\{ \sum_{k=1}^{\infty} |m_{k,n+1}|, \sum_{k=1}^{\infty} |m_{k,n+2}|, \dots \right\} \\ &\leq C_1 \sup\{r^{n+1}, r^{n+2}, \dots\} \\ &= C_1r^{n+1} \rightarrow 0. \end{aligned}$$

Thus,  $M_{r,\infty}$  is compact as desired. □

**COROLLARY 1.**  $M_{\infty}$  is compact on  $l^1 \times l^1$ , where for  $\underline{x} = (\underline{x}^1, \underline{x}^2) \in l^1 \times l^1$ ,

$$(12) \quad M_{\infty} \begin{pmatrix} \underline{x}^1 \\ \underline{x}^2 \end{pmatrix} = \begin{pmatrix} \sum_{k=1}^{\infty} \sqrt{kj}\eta_{k+j-1}x^1_k + \sum_{k=1}^{\infty} \sqrt{kj}\gamma_{k+j-1}x^2_k \\ \sum_{k=1}^{\infty} \sqrt{kj}\gamma_{k+j-1}x^1_k - \sum_{k=1}^{\infty} \sqrt{kj}\eta_{k+j-1}x^2_k \end{pmatrix}.$$

The norm on  $l^1 \times l^1$  is given by

$$\|\underline{x}\|_{l^1 \times l^1} = \|\underline{x}^1\|_{l^1} + \|\underline{x}^2\|_{l^1}.$$

*Proof.* From the notation of the problem, it is easily verified that

$$\|M_\infty - M_n\|_{l^1 \times l^1} \leq \|M_{r,\infty} - M_{r,n}\|_{l^1} + \|M_{i,\infty} - M_{i,n}\|_{l^1}.$$

The result follows from Theorem 1.  $\square$

Next, we discuss the discretization of (10). Since  $M_\infty$  is compact and the matrix-vector multiplications can be performed rapidly, we will solve the discrete (normal) equations using the conjugate gradient method on the subspace orthogonal to the one-dimensional null space.

**4. Discretization.** The natural choice for discretization is to truncate the sums given in (12) to  $n$ . This will lead to finite linear systems. However, in practice one does not have the exact Fourier coefficients. If the conformal map  $f$  is known explicitly, we approximate the  $h_k$ 's by evaluating  $d(z) := f(z)/\overline{f'(z)}$  at the  $N = 2n$  Fourier points,  $z = e^{ij\pi/n}, j = 0, 1, \dots, N - 1$ , and taking the  $N$ -point FFT. In this case, the discrete  $h_1, \dots, h_n$  decay at a similar rate to the exact  $h_k$  (see [He, eq. 13.2-8, p. 20]). However, since the discrete Fourier coefficients are  $N$ -periodic,  $h_k = h_{k-N}$ , the remaining coefficients  $h_{n+1} = h_{-n+1}, \dots, h_{N-1} = h_{-1}$  do not decay geometrically. We just set  $h_k = 0, k > n$ , to insure geometric decay. When  $f$  is not known exactly, we use a numerical approximation at the  $N$  Fourier points given by Wegmann's method, as discussed in section 5, and again set  $h_k = 0$  for  $k = n + 1, \dots, N - 1$ . To avoid introducing more notation, we now let  $h_k, A_k$ , etc., denote the discrete Fourier coefficients.

The notation is similar to the infinite-dimensional case

$$\begin{aligned} D_n &= \text{diag}(1, 2, \dots, n), \\ \underline{\alpha} &= (\text{Re } a_1, \dots, \text{Re } a_n)^T, \quad \underline{\beta} = (\text{Im } a_1, \dots, \text{Im } a_n)^T, \\ \underline{B} &= (\text{Re } A_1, \dots, \text{Re } A_n)^T, \quad \underline{C} = (\text{Im } A_1, \dots, \text{Im } A_n)^T, \\ \underline{x} &= \begin{pmatrix} D_n^{1/2} \underline{\alpha} \\ D_n^{1/2} \underline{\beta} \end{pmatrix}, \quad \underline{r} = \begin{pmatrix} D_n^{1/2} \underline{B} \\ D_n^{1/2} \underline{C} \end{pmatrix}, \end{aligned}$$

and

$$H_n = \begin{pmatrix} H_{r,n} & H_{i,n} \\ H_{i,n} & -H_{r,n} \end{pmatrix}.$$

Then, analogously to the infinite system, we have

$$M_n = \begin{pmatrix} M_{r,n} & M_{i,n} \\ M_{i,n} & -M_{r,n} \end{pmatrix} = \begin{pmatrix} D_n^{1/2} H_{r,n} D_n^{1/2} & D_n^{1/2} H_{i,n} D_n^{1/2} \\ D_n^{1/2} H_{i,n} D_n^{1/2} & -D_n^{1/2} H_{r,n} D_n^{1/2} \end{pmatrix},$$

so that our problem is to solve

$$(13) \quad (I_n + M_n)\underline{x} = \underline{r}.$$

Recall that  $\underline{x}$  is subject to a uniqueness condition. Since  $f'(0) > 0$ , the condition  $\text{Im}(a_1/f'(0)) = 0$  implies  $x_{n+1} = 0$ . Clearly, the  $(k, j)$ th entry of  $M_{r,n}$  and  $M_{i,n}$  are, respectively,  $\sqrt{kj}\text{Re}(h_{k+j})$  and  $\sqrt{kj}\text{Im}(h_{k+j})$ .

We have computed the eigenvalues of  $M_n$  for the examples in section 5 using MATLAB. Note that if  $\mu$  is an eigenvalue of  $M_n$ , then  $-\mu$  is also an eigenvalue. We also find that  $-1$  is an eigenvalue of  $M_n$ . The rest of the eigenvalues decay rapidly to 0. The decay is due to the compactness of  $M_\infty$  shown in Corollary 1 of section 3. By [An], the spectrum of  $M_n$  is near to the spectrum of  $M_\infty$  for large  $n$ . We solve the normal equations by conjugate gradient, since  $(I_n + M_n)^2$  is positive semidefinite. In our examples, so far, we have noticed that  $I_n + M_n$  is also positive semidefinite, but we have no proof of this, in general. In these examples, we have used conjugate gradient directly on (13) with some computational savings. (13) could also no doubt be solved efficiently with MINRES. In addition, we have solved (a truncated version) of the nonsymmetric system (9) with GMRES with very good results. We hope to address these issues further in future work.

Recall that our infinite system (10) has a one-dimensional null space. The null space is generated by the null vector,

$$\underline{v} = (-\text{Im } c_1, -\sqrt{2}\text{Im } c_2, \dots, -\sqrt{k}\text{Im } c_k, \dots, \text{Re } c_1, \sqrt{2}\text{Re } c_2, \dots, \sqrt{k}\text{Re } c_k, \dots)^T.$$

In the discrete case, we find that for large  $n$ ,

$$\underline{v} = (-\text{Im } c_1, -\sqrt{2}\text{Im } c_2, \dots, -\sqrt{n}\text{Im } c_n, \text{Re } c_1, \sqrt{2}\text{Re } c_2, \dots, \sqrt{n}\text{Re } c_n)^T$$

satisfies  $(I_n + M_n)\underline{v} = \underline{0}$  to within discretization error using our discrete approximations to the  $c_k$ 's. It follows that our solution can be decomposed as

$$\underline{y} = \underline{x} + \delta \underline{v}.$$

It is clear from the conjugate gradient algorithm that if the initial guess  $\underline{x}^{(0)}$  is in  $\underline{v}^\perp$  then subsequent iterates  $\underline{x}^{(q)}$  will be in  $\underline{v}^\perp$ . We take  $\underline{x}^{(0)} = \underline{0}$ . Conjugate gradient will then find  $\underline{x} \in \underline{v}^\perp$  and, imposing the uniqueness condition,  $y_{n+1} = 0$  will give us  $\delta$ . By the results above,  $(I_n + M_n)^2$  restricted to  $\underline{v}^\perp$  is positive definite for sufficiently large  $n$ , since the second smallest eigenvalue of  $I_\infty + M_\infty$  is bounded away from 0. Therefore, conjugate gradient can be applied to the normal equations and the method will converge superlinearly.

In addition, we note that the matrix-vector multiplication involving the matrix  $M_n$  can be done efficiently using FFTs. In fact, for any  $n$ -vector  $\underline{y}$ , since  $D_n$  is diagonal,  $D_n^{1/2}\underline{y}$  can be computed in  $n$  operations. Moreover, the matrix-vector multiplication  $H\underline{y}$ , where  $H$  is the Hankel matrix  $H_{r,n}$  or  $H_{i,n}$ , can be computed in  $O(N \log N)$  by using FFTs. The idea is to compute  $T\underline{s} = (HJ)(J\underline{y})$  where  $J$  is the reversion matrix with 1's on the anti-diagonal, and  $T$  is a Toeplitz matrix (constant along diagonals). Next we imbed  $T$  into a matrix  $C$  as follows:

$$C = \begin{pmatrix} T & X \\ X & T \end{pmatrix},$$

where  $X$  is chosen to make  $C$  circulant. Now  $C$  can be decomposed as  $C = F^* \Lambda F$ , where  $F$  is the  $N$ -point Fourier matrix and  $\Lambda$  is a diagonal matrix containing the eigenvalues of  $C$ . For more details on fast methods for Hankel and Toeplitz matrices see, e.g., [CN].

**5. Numerical examples.** In examples (i), (ii), and (iii), we choose  $\phi(\zeta) = \zeta^3$  and  $\chi(\zeta) = 0$ . Then  $u(\eta, \mu) = \eta^4 - \mu^4$ . Note that, for the conformal map  $f(z)$  from the disk,  $\phi(z) = (f(z))^3$  and the boundary values at the mesh points are given by



$G(z) = 4(\operatorname{Re}f(z))^3 - i4(\operatorname{Im}f(z))^3$ . The discretization error in the tables is given by the sup norm

$$\max_{0 \leq j \leq N-1} |\phi(e^{i2\pi j/N}) - \phi_n(e^{i2\pi j/N})|,$$

where  $\phi_n$  is our  $n$ th-degree approximation to  $\phi$ . For analytic curves, this error behaves similarly to the discretization error for the conformal map, which is  $O(R^N)$ , with  $R$  as given in Lemma 1; see [De] for a discussion of the accuracy of the conformal mapping methods.

We use the FFT method in [Weg] to find the approximate conformal map  $f$ . Wegmann approximates  $f$  by solving a discrete interpolation problem on the unit disk: find  $P_{n+1}(z)$ , a polynomial of degree  $n+1$ , such that  $P_{n+1}(e^{i2\pi j/N}) \in \Gamma, j = 0, \dots, N-1$ , with the normalization that the  $P_{n+1}(0)$  is fixed and the coefficients of  $z$  and  $z^{n+1}$  are real. He computes this polynomial by applying a Newton method to find a discrete approximation to the boundary correspondence. The linear systems may be solved by the conjugate gradient method in  $O(N \log N)$  per step. Quadratic convergence of the Newton iterations and convergence of the polynomial to the conformal map as  $N$  increases for sufficiently smooth  $\Gamma$  is proven. Numerical experiments indicate that this method is among the most robust and reliable of the Fourier series methods on the disk [De].

For examples where the exact  $f$  is known,  $d = f/\overline{f'}$  may be computed with either the exact or the approximate  $f$ . This seems to make little difference in the calculations if the approximate  $f$  is sufficiently accurate. The timings for finding the approximate  $f$  using [Weg] are usually only slightly greater than the timings given in the tables for solving the boundary value problem for a given  $N$ .

In the tables below, iter is the number of iterations required by conjugate gradient for the residuals to be  $\leq 10^{-14}$ . The computations were done in double precision on the WSU IBM ES9121 Model 440 mainframe computer, and some rough timings are given. (Figures 1 and 2 and some of our examples were also done in MATLAB, with similar results.) Stopping the iterations after the level of discretization error has been achieved could further reduce the timings, though not dramatically for these examples of very fast superlinear convergence. Note that as the minor-to-major axis ratio  $\alpha$  of a region decreases toward 0 (that is, as  $R$  in Lemma 1 increases to 1), the convergence rate of the conjugate gradient method decreases. In our examples below,  $R$  may be taken as the distance from the origin to the nearest singularity of  $f$ , and the connection with the minor-to-major aspect ratio is known [De]. In a future paper we will show how the convergence rate of the conjugate gradient method depends on the smoothness of the boundary  $\Gamma$ .

Other cases were also tried successfully, such as the simple examples in [KK]. If the biharmonic function has too simple a Goursat representation, the iterations may converge artificially fast. For instance, if  $u(\eta, \mu) = \eta^2 + \eta\mu + \mu^2$ , then  $\phi(z) = f(z)$  and convergence is achieved in one iteration if  $N$  is large enough. On the other hand, note that the 5-to-1 ellipse in [GGMa, Table 3] is a difficult region for the conformal map from the disk and would require large  $N$ . See example (iii) below.

*Example (i), inverted ellipse.*  $\Gamma : \gamma(\sigma) = \rho(\sigma)e^{i\sigma}$ , where  $\rho(\sigma) = \sqrt{1 - (1 - \alpha^2)\sin^2 \sigma}$  for  $0 \leq \sigma \leq 2\pi$  and  $0 < \alpha \leq 1$ . This map is derived by inverting the familiar Joukowski map to the exterior of an ellipse. We have

$$f(z) = \frac{2\alpha z}{1 + \alpha - (1 - \alpha)z^2}.$$

TABLE 1  
*Inverted ellipse with exact map and conjugate gradient.*

$\alpha$	$N$	Discr. error	iter	CPU sec
.8	32	$.5 \cdot 10^{-5}$	4	.2
.8	64	$.5 \cdot 10^{-12}$	3	.2
.8	128	$.5 \cdot 10^{-14}$	3	.2
.4	64	$.1 \cdot 10^{-3}$	6	.2
.4	128	$.6 \cdot 10^{-9}$	4	.2
.4	256	$.1 \cdot 10^{-13}$	4	.3
.2	128	$.2 \cdot 10^{-3}$	6	.3
.2	256	$.2 \cdot 10^{-8}$	4	.3
.2	512	$.3 \cdot 10^{-13}$	4	.4




FIG. 1. *Eigenvalue distribution for  $(I_n + M_n)^2$  for the arctanh region (example (ii)) with  $\alpha = .49$  and  $N = 256$ .*

See Table 1 for results. Notice how iter is roughly independent of  $N$ , but increases as  $\alpha \downarrow 0$  in our examples.

*Example (ii), arctanh.* Here the conformal map is given by  $f(z) = \log((1+rz)/(1-rz))$ ,  $0 < r < 1$ , which maps the disk to increasingly elongated, cigar-shaped regions as  $r \uparrow 1$ . This map is perhaps the simplest example of a conformal map exhibiting the exponential crowding [De]. Figure 1 shows roughly seven outlying eigenvalues of  $(I_n + M_n)^2$  for  $\alpha = .49$ . Thus conjugate gradient for the normal equations takes about seven iterations to converge, as one would expect; see Table 2 and Fig. 2. Also note the semilog plot in Figure 2 that shows the superlinear convergence behavior of the residuals (using MATLAB).

TABLE 2  
*Arctanh regions with exact map and conjugate gradient.*

$\alpha(r)$	$N$	Discr. error	iter	CPU sec
.84 (.5)	32	$.4 \cdot 10^{-4}$	6	.2
.84 (.5)	64	$.4 \cdot 10^{-9}$	4	.2
.84 (.5)	128	$.8 \cdot 10^{-14}$	4	.2
.49 (.9)	128	$.6 \cdot 10^{-3}$	8	.4
.49 (.9)	256	$.4 \cdot 10^{-6}$	6	.3
.49 (.9)	512	$.4 \cdot 10^{-12}$	6	.5
.29 (.99)	512	$.4 \cdot 10^{-1}$	14	.9
.29 (.99)	1024	$.2 \cdot 10^{-2}$	13	1.5
.29 (.99)	2048	$.6 \cdot 10^{-5}$	12	2.8
.29 (.99)	4096	$.1 \cdot 10^{-9}$	12	6.0




FIG. 2. *Convergence of residuals for seven iterations of the conjugate gradient method for the normal equations for the arctanh region (example (ii)) with  $\alpha = .49$  and  $N = 256$ .*

*Example (iii), ellipse.* Here  $\Gamma : \gamma(\sigma) = \rho(\sigma)e^{i\sigma}$ , where  $\rho(\sigma) = \alpha/\sqrt{1 - (1 - \alpha^2)\cos^2\sigma}$  for  $0 \leq \sigma \leq 2\pi$  and  $0 < \alpha \leq 1$ . The exact map can be given in terms of an elliptic integral. This case also exhibits exponential crowding [De]. We approximate the  $f$  with [Weg]. See Table 3.

**Acknowledgment.** The authors thank Rudolf Wegmann for providing a copy of his Fortran code based on [Weg].

TABLE 3  
*Ellipses with approximate map and conjugate gradient.*

$\alpha$	$N$	Discr. error	iter	CPU sec
.8	32	$.7 \cdot 10^{-3}$	6	.2
.8	64	$.5 \cdot 10^{-6}$	4	.2
.8	128	$.4 \cdot 10^{-12}$	3	.2
.8	256	$.2 \cdot 10^{-13}$	3	.3
.6	128	$.6 \cdot 10^{-3}$	8	.3
.6	256	$.3 \cdot 10^{-6}$	6	.4
.6	512	$.2 \cdot 10^{-12}$	3	.4
.4	2048	$.2 \cdot 10^{-4}$	10	2.6
.4	4096	$.7 \cdot 10^{-10}$	8	4.6

## REFERENCES

- [An] P. M. ANSELONE, *Collectively Compact Operator Approximation Theory*, Prentice-Hall, Englewood Cliffs, NJ, 1971.
- [CN] R. H. CHAN AND M. K. NG, *Conjugate gradient methods for Toeplitz systems*, SIAM Rev., 38 (1996), pp. 427–482.
- [Con] J. B. CONWAY, *A Course in Functional Analysis*, 2nd ed., Springer-Verlag, Berlin, 1990.
- [De] T. K. DELILLO, *The accuracy of numerical conformal mapping methods: A survey of examples and results*, SIAM J. Numer. Anal., 31 (1994), pp. 788–812.
- [DE] T. K. DELILLO AND A. R. ELCRAT, *A Fornberg-like conformal mapping method for slender regions*, J. Comput. Appl. Math., 46 (1993), pp. 49–64.
- [DEP] T. K. DELILLO, A. R. ELCRAT, AND J. A. PFALTZGRAFF, *Numerical Conformal Mapping Methods Based on Faber Series*, preprint.
- [Ga] D. GAIER, *Konstruktive Methoden der konformen Abbildung*, Springer-Verlag, Berlin, 1964.
- [GVL] G. GOLUB AND C. VAN LOAN, *Matrix Computations*, 2nd ed., The Johns Hopkins University Press, Baltimore, MD, 1989.
- [GGMa] A. GREENBAUM, L. GREENGARD, AND A. MAYO, *On the numerical solution of the bi-harmonic equation in the plane*, Phys. D, 60 (1992), pp. 216–225.
- [He] P. HENRICI, *Applied and Computational Complex Analysis, Vol. III*, John Wiley, New York, 1986.
- [KK] L. V. KANTOROVICH AND V. I. KRYLOV, *Approximate Methods of Higher Analysis*, P. Noordhoff Groningen, the Netherlands, 1958.
- [K] A. KARAGEORGHIS, *The method of fundamental solutions for the solution of steady-state free boundary problems*, J. Comput. Phys., 98 (1992), pp. 119–128.
- [MG] A. MAYO AND A. GREENBAUM, *Fast parallel iterative solution of Poisson's and the bi-harmonic equation on irregular regions*, SIAM J. Sci. Statist. Comput., 13 (1992), pp. 101–118.
- [Mik] S. G. MIKLIN, *Integral Equations*, MacMillan, New York, 1964.
- [MT] L. M. MILNE-THOMSON, *Theoretical Hydrodynamics*, 5th ed., MacMillan, London, 1968.
- [Musk] N. I. MUSKHELISHVILI, *Some Basic Problems of the Mathematical Theory of Elasticity*, P. Noordhoff Groningen, the Netherlands, 1953.
- [Poz] C. POZRIKIDIS, *Boundary Integral and Singularity Methods for Linearized Viscous Flow*, Cambridge University Press, UK, 1992.
- [Weg] R. WEGMANN, *Discrete Riemann-Hilbert problems, interpolation of simply closed curves, and numerical conformal mapping*, J. Comput. Appl. Math., 23 (1988), pp. 323–352.