

Section 1.5

2.) $f(x) = x^2 - 3x + 1$

x	$f(x)$
1.9	-1.09
1.99	-1.0099
1.999	-1.000999
2.001	-0.998999
2.01	-0.9899
2.1	-0.89

Guess: $\lim_{x \rightarrow 2} (x^2 - 3x + 1) = -1$

6.) $f(x) = \frac{\sqrt{x+2} - \sqrt{2}}{x}$

x	$f(x)$
-0.1	0.358086
-0.01	0.353996
-0.001	0.353597
0.001	0.353509
0.01	0.353112
0.1	0.349241

Guess: $\lim_{x \rightarrow 0} \frac{\sqrt{x+2} - \sqrt{2}}{x} \approx 0.3535$

9.) a.) $\lim_{x \rightarrow 0} f(x) = 1$

b.) $\lim_{x \rightarrow -1} f(x) = 3$

10.) a.) $\lim_{x \rightarrow 1} f(x) = -2$

b.) $\lim_{x \rightarrow 3} f(x) = 0$

11.) a.) $\lim_{x \rightarrow 0} g(x) = 1$

b.) $\lim_{x \rightarrow -1} g(x) = 3$

12.) a.) $\lim_{x \rightarrow -2} h(x) = -5$

b.) $\lim_{x \rightarrow 0} h(x) = -3$

17.) a.) $\lim_{x \rightarrow 3^+} f(x) = 1$

b.) $\lim_{x \rightarrow 3^-} f(x) = 1$

c.) $\lim_{x \rightarrow 3} f(x) = 1$

19.) a.) $\lim_{x \rightarrow 3^+} f(x) = 0$

b.) $\lim_{x \rightarrow 3^-} f(x) = 0$

c.) $\lim_{x \rightarrow 3} f(x) = 0$

$$22.) \text{ a.) } \lim_{x \rightarrow -1^+} f(x) = 0$$

$$\text{ b.) } \lim_{x \rightarrow -1^-} f(x) = 2$$

$$\text{ c.) } \lim_{x \rightarrow -1} f(x) \text{ DNE}$$

$$23.) \lim_{x \rightarrow 2} x^4 = 2^4 = 16$$

$$29.) \lim_{x \rightarrow 3} \sqrt{x+1} = \sqrt{4} = 2$$

$$32.) \lim_{x \rightarrow -2} \frac{3x+1}{2-x} = \frac{-5}{4}$$

$$37.) \lim_{x \rightarrow 3} \frac{\sqrt{x+1}-1}{x} = \frac{2-1}{3} = \frac{1}{3}$$

$$40.) \lim_{x \rightarrow 2} \frac{\frac{1}{x+2} - \frac{1}{2}}{x} = \frac{\frac{1}{4} - \frac{1}{2}}{2} = \frac{-\frac{1}{4}}{2} = -\frac{1}{8}$$

$$42.) \lim_{x \rightarrow -1} \frac{2x^2 - x - 3}{x+1} \stackrel{\frac{0}{0}}{=} \lim_{x \rightarrow -1} \frac{(2x-3)(x+1)}{\cancel{x+1}}$$
$$= \lim_{x \rightarrow -1} (2x-3) = -5$$

$$44.) \lim_{x \rightarrow 2} \frac{2-x}{x^2-4} \stackrel{\frac{0}{0}}{=} \lim_{x \rightarrow 2} \frac{-(x-2)}{(x-2)(x+2)} = \frac{-1}{4}$$

$$47.) \lim_{x \rightarrow -2} \frac{x^3+8}{x+2} \stackrel{\frac{0}{0}}{=} \lim_{x \rightarrow -2} \frac{(x+2)(x^2-2x+4)}{\cancel{x+2}}$$
$$= 4+4+4 = 12$$


50.) since

$$\lim_{x \rightarrow 2^+} \frac{|x-2|}{x-2} = \lim_{x \rightarrow 2^+} 1 = 1$$

and

$$\lim_{x \rightarrow 2^-} \frac{|x-2|}{x-2} = \lim_{x \rightarrow 2^-} -1 = -1$$

so $\lim_{x \rightarrow 2} \frac{|x-2|}{x-2}$ does not exist.




51.)

$$f(x) = \begin{cases} \frac{1}{3}x - 2, & x \leq 3 \\ -2x + 5, & x > 3 \end{cases}$$

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (-2x + 5) = -1$$

$$\text{and } \lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} \left(\frac{1}{3}x - 2\right) = -1$$

$$\text{so } \lim_{x \rightarrow 3} f(x) = -1.$$



52.) $f(s) = \begin{cases} s, & s \leq 1 \\ 1-s, & s > 1 \end{cases}$ so

$$\lim_{s \rightarrow 1^+} f(s) = \lim_{s \rightarrow 1^+} (1-s) = 1 - (1) = 0 \quad \text{and}$$

$$\lim_{s \rightarrow 1^-} f(s) = \lim_{s \rightarrow 1^-} s = 1, \text{ so } \lim_{s \rightarrow 1} f(s) \text{ does not exist.}$$

56.) $\lim_{\Delta x \rightarrow 0} \frac{\sqrt{x+\Delta x} - \sqrt{x}}{\Delta x} = \frac{0}{0}$

$$= \lim_{\Delta x \rightarrow 0} \frac{\sqrt{x+\Delta x} - \sqrt{x}}{\Delta x} \cdot \frac{\sqrt{x+\Delta x} + \sqrt{x}}{\sqrt{x+\Delta x} + \sqrt{x}}$$

$$= \lim_{\Delta X \rightarrow 0} \frac{(X+\Delta X) - X}{\Delta X (\sqrt{X+\Delta X} + \sqrt{X})} = \lim_{\Delta X \rightarrow 0} \frac{\Delta X}{\Delta X (\sqrt{X+\Delta X} + \sqrt{X})}$$

$$= \frac{1}{\sqrt{X} + \sqrt{X}} = \frac{1}{2\sqrt{X}}$$

57.) $\lim_{\Delta t \rightarrow 0} \frac{(t+\Delta t)^2 - 5(t+\Delta t) - (t^2 - 5t)}{\Delta t}$


"0/0"
 $= \lim_{\Delta t \rightarrow 0} \frac{t^2 + 2t \cdot \Delta t + (\Delta t)^2 - 5t - 5 \cdot \Delta t - t^2 + 5t}{\Delta t}$

$$= \lim_{\Delta t \rightarrow 0} \frac{\Delta t \cdot (2t + \Delta t - 5)}{\Delta t} = 2t + 0 - 5 = 2t - 5$$

59.) $\lim_{x \rightarrow 1^-} \frac{2}{x^2 - 1} = \frac{"2"}{0^-} = -\infty$




60.) $\lim_{x \rightarrow 1^+} \frac{5}{1-x} = \frac{"5"}{0^-} = -\infty$




61.) $\lim_{x \rightarrow -2^-} \frac{1}{x+2} = \frac{"1"}{0^-} = -\infty$

62.) $\lim_{x \rightarrow 0^-} \frac{x+1}{x} = \frac{"1"}{0^-} = -\infty$



$$64.) \quad \lim_{x \rightarrow 1} \frac{x^2 + 6x - 7}{x^3 - x^2 + 2x - 2} \stackrel{0/0}{=} \lim_{x \rightarrow 1} \frac{\cancel{(x-1)}(x+7)}{\cancel{(x-1)}(x^2+2)} \\ = \lim_{x \rightarrow 1} \frac{x+7}{x^2+2} = \frac{8}{3}$$

$$\begin{array}{r} x^2 + 2 \\ x-1 \overline{) x^3 - x^2 + 2x - 2} \\ \underline{x^3 - x^2} \\ 2x - 2 \\ \underline{2x - 2} \\ 0 \end{array}$$



Since $4 - x^2 \leq f(x) \leq 4 + x^2$ it follows that

$$\lim_{x \rightarrow 0} (4 - x^2) \leq \lim_{x \rightarrow 0} f(x) \leq \lim_{x \rightarrow 0} (4 + x^2)$$

or

$$4 \leq \lim_{x \rightarrow 0} f(x) \leq 4 ;$$

thus


$$\lim_{x \rightarrow 1} f(x) = 4 .$$

Worksheet 1

11.) a.) $Y = \frac{2-3x}{x-1} \rightarrow Y(x-1) = 2-3x \rightarrow$
 $XY - Y = 2-3x \rightarrow XY + 3x = Y+2 \rightarrow$
 $x(Y+3) = Y+2 \rightarrow x = \frac{Y+2}{Y+3} = f^{-1}(Y)$

(or $Y = \frac{x+2}{x+3} = f^{-1}(x)$)

b.) $Y = x^2 - 4x$ for $x \leq 2 \rightarrow$
 $Y = (x^2 - 4x + 4) - 4$ for $x \leq 2 \rightarrow$
 $Y = (x-2)^2 - 4 \rightarrow Y+4 = (x-2)^2 \rightarrow$



why? $x-2 = \pm\sqrt{Y+4} \rightarrow x = 2 \pm \sqrt{Y+4} \rightarrow$

$x = 2 - \sqrt{Y+4} = f^{-1}(Y)$

(or $Y = 2 - \sqrt{x+4} = f^{-1}(x)$)



12.) $f(g(x)) = \frac{g(x)}{g(x)+5} = 5-x^3 \rightarrow g(x) = (5-x^3)(g(x)+5) \rightarrow$

$g(x) = 5g(x) - x^3g(x) + 25 - 5x^3 \rightarrow$

$x^3g(x) - 4g(x) = 25 - 5x^3 \rightarrow (x^3 - 4)g(x) = 25 - 5x^3 \rightarrow$


$g(x) = \frac{25 - 5x^3}{x^3 - 4}$

14.) a.) Use similar triangles:




$\frac{5}{4} = \frac{x}{3} \rightarrow x = \frac{15}{4}$

b.)



Use similar triangles:



$$\frac{6}{8} = \frac{x}{y} \rightarrow y = \frac{4}{3}x \quad \text{and}$$

$$\frac{6}{8} = \frac{10-y}{x} \rightarrow \frac{3}{4}x = 10-y \rightarrow \frac{3}{4}x = 10 - \frac{4}{3}x \rightarrow$$

$$\left(\frac{3}{4} + \frac{4}{3}\right)x = 10 \rightarrow \frac{25}{12}x = 10 \rightarrow x = \frac{24}{5}$$

15.) a.) IV b.) II c.) III

16.) heart rate




19.) domain: $-1 \leq x \leq 1.5$

range: $-1.5 \leq y \leq 0$

23.) Let x : miles traveled

a.) Co. A: cost $C = 40 + 0.15x = A(x)$

Co. B: cost $C = 50 + 0.1x = B(x)$



Find pt. of intersection:

$$40 + 0.15x = 50 + 0.1x \rightarrow$$

$$0.05x = 10 \rightarrow$$

$$x = 200 \text{ mi.}$$


c.) Use Co. A if $x \leq 200$ miles;
use Co. B if $x > 200$ miles.

SA2: Volume $V = \frac{4}{3}\pi r^3 = \frac{\pi}{6} \rightarrow r^3 = \frac{1}{8} \rightarrow$
 $r = \frac{1}{2}$ ft. so diameter of balls is $d = 1$ ft.

a.) The number of balls will be
 $6 \times 4 \times 3 = 72$ balls

b.) The available space in box is
(Volume of box) - (Volume of balls)
 $= 72 - 72\left(\frac{\pi}{6}\right) \approx 34.3$ ft.³, so total weight
of H₂O is $W = (34.3)(62.5) \approx 2143.81$ lbs.

SA5:



Circumference
 $C = 2\pi r = 2\pi \rightarrow r = 1$ ft.,
by Pythagorean Theorem
 $x^2 + x^2 = 2^2 \rightarrow$

$$2x^2 = 4 \rightarrow x = \sqrt{2} \text{ ft.}$$

so perimeter of square is

$$P = 4\sqrt{2} \text{ ft.}$$