

FIGURE 16.36 The region bounded by the circle  $C_h$  and the curve  $K$ .

which leads to the conclusion that

$$\oint_K (M dx + N dy) = 2\pi$$

for any such curve  $K$ . We can explain this result by changing to polar coordinates. With

$$\begin{aligned} x &= r \cos \theta, & y &= r \sin \theta, \\ dx &= -r \sin \theta d\theta + \cos \theta dr, & dy &= r \cos \theta d\theta + \sin \theta dr, \end{aligned}$$

we have

$$\frac{x dy - y dx}{x^2 + y^2} = \frac{r^2(\cos^2 \theta + \sin^2 \theta) d\theta}{r^2} = d\theta,$$

and  $\theta$  increases by  $2\pi$  as we traverse  $K$  once counterclockwise.

## EXERCISES 16.4

### Verifying Green's Theorem

In Exercises 1–4, verify the conclusion of Green's Theorem by evaluating both sides of Equations (3) and (4) for the field  $\mathbf{F} = M\mathbf{i} + N\mathbf{j}$ . Take the domains of integration in each case to be the disk  $R: x^2 + y^2 \leq a^2$  and its bounding circle  $C: \mathbf{r} = (a \cos t)\mathbf{i} + (a \sin t)\mathbf{j}$ ,  $0 \leq t \leq 2\pi$ .

- $\mathbf{F} = -y\mathbf{i} + x\mathbf{j}$
- $\mathbf{F} = y\mathbf{i}$
- $\mathbf{F} = 2x\mathbf{i} - 3y\mathbf{j}$
- $\mathbf{F} = -x^2y\mathbf{i} + xy^2\mathbf{j}$

### Counterclockwise Circulation and Outward Flux

In Exercises 5–10, use Green's Theorem to find the counterclockwise circulation and outward flux for the field  $\mathbf{F}$  and curve  $C$ .

- $\mathbf{F} = (x - y)\mathbf{i} + (y - x)\mathbf{j}$   
 C: The square bounded by  $x = 0, x = 1, y = 0, y = 1$
- $\mathbf{F} = (x^2 + 4y)\mathbf{i} + (x + y^2)\mathbf{j}$   
 C: The square bounded by  $x = 0, x = 1, y = 0, y = 1$
- $\mathbf{F} = (y^2 - x^2)\mathbf{i} + (x^2 + y^2)\mathbf{j}$   
 C: The triangle bounded by  $y = 0, x = 3$ , and  $y = x$
- $\mathbf{F} = (x + y)\mathbf{i} - (x^2 + y^2)\mathbf{j}$   
 C: The triangle bounded by  $y = 0, x = 1$ , and  $y = x$
- $\mathbf{F} = (x + e^x \sin y)\mathbf{i} + (x + e^x \cos y)\mathbf{j}$   
 C: The right-hand loop of the lemniscate  $r^2 = \cos 2\theta$
- $\mathbf{F} = \left(\tan^{-1} \frac{y}{x}\right)\mathbf{i} + \ln(x^2 + y^2)\mathbf{j}$   
 C: The boundary of the region defined by the polar coordinate inequalities  $1 \leq r \leq 2, 0 \leq \theta \leq \pi$
- Find the counterclockwise circulation and outward flux of the field  $\mathbf{F} = xy\mathbf{i} + y^2\mathbf{j}$  around and over the boundary of the region enclosed by the curves  $y = x^2$  and  $y = x$  in the first quadrant.

- Find the counterclockwise circulation and the outward flux of the field  $\mathbf{F} = (-\sin y)\mathbf{i} + (x \cos y)\mathbf{j}$  around and over the square cut from the first quadrant by the lines  $x = \pi/2$  and  $y = \pi/2$ .
- Find the outward flux of the field

$$\mathbf{F} = \left(3xy - \frac{x}{1 + y^2}\right)\mathbf{i} + (e^x + \tan^{-1} y)\mathbf{j}$$

across the cardioid  $r = a(1 + \cos \theta)$ ,  $a > 0$ .

- Find the counterclockwise circulation of  $\mathbf{F} = (y + e^x \ln y)\mathbf{i} + (e^x/y)\mathbf{j}$  around the boundary of the region that is bounded above by the curve  $y = 3 - x^2$  and below by the curve  $y = x^4 + 1$ .

### Work

In Exercises 15 and 16, find the work done by  $\mathbf{F}$  in moving a particle once counterclockwise around the given curve.

- $\mathbf{F} = 2xy^3\mathbf{i} + 4x^2y^2\mathbf{j}$   
 C: The boundary of the "triangular" region in the first quadrant enclosed by the  $x$ -axis, the line  $x = 1$ , and the curve  $y = x^3$
- $\mathbf{F} = (4x - 2y)\mathbf{i} + (2x - 4y)\mathbf{j}$   
 C: The circle  $(x - 2)^2 + (y - 2)^2 = 4$

### Evaluating Line Integrals in the Plane

Apply Green's Theorem to evaluate the integrals in Exercises 17–20.

- $\oint_C (y^2 dx + x^2 dy)$   
 C: The triangle bounded by  $x = 0, x + y = 1, y = 0$
- $\oint_C (3y dx + 2x dy)$   
 C: The boundary of  $0 \leq x \leq \pi, 0 \leq y \leq \sin x$

19. 
$$\oint_C (6y + x) dx + (y + 2x) dy$$

$C$ : The circle  $(x - 2)^2 + (y - 3)^2 = 4$

20. 
$$\oint_C (2x + y^2) dx + (2xy + 3y) dy$$

$C$ : Any simple closed curve in the plane for which Green's Theorem holds

### Calculating Area with Green's Theorem

If a simple closed curve  $C$  in the plane and the region  $R$  it encloses satisfy the hypotheses of Green's Theorem, the area of  $R$  is given by

#### Green's Theorem Area Formula

$$\text{Area of } R = \frac{1}{2} \oint_C x dy - y dx \quad (13)$$

The reason is that by Equation (3), run backward,

$$\begin{aligned} \text{Area of } R &= \iint_R dy dx = \iint_R \left( \frac{1}{2} + \frac{1}{2} \right) dy dx \\ &= \oint_C \frac{1}{2} x dy - \frac{1}{2} y dx. \end{aligned}$$


Use the Green's Theorem area formula (Equation 13) to find the areas of the regions enclosed by the curves in Exercises 21–24.

21. The circle  $\mathbf{r}(t) = (a \cos t)\mathbf{i} + (a \sin t)\mathbf{j}$ ,  $0 \leq t \leq 2\pi$

22. The ellipse  $\mathbf{r}(t) = (a \cos t)\mathbf{i} + (b \sin t)\mathbf{j}$ ,  $0 \leq t \leq 2\pi$

23. The astroid  $\mathbf{r}(t) = (\cos^3 t)\mathbf{i} + (\sin^3 t)\mathbf{j}$ ,  $0 \leq t \leq 2\pi$

24. The curve  $\mathbf{r}(t) = t^2\mathbf{i} + ((t^3/3) - t)\mathbf{j}$ ,  $-\sqrt{3} \leq t \leq \sqrt{3}$  (see accompanying figure).



### Theory and Examples

25. Let  $C$  be the boundary of a region on which Green's Theorem holds. Use Green's Theorem to calculate

a. 
$$\oint_C f(x) dx + g(y) dy$$

b. 
$$\oint_C ky dx + hx dy \quad (k \text{ and } h \text{ constants}).$$

26. **Integral dependent only on area** Show that the value of

$$\oint_C xy^2 dx + (x^2y + 2x) dy$$

around any square depends only on the area of the square and not on its location in the plane.

27. What is special about the integral

$$\oint_C 4x^3y dx + x^4 dy?$$

Give reasons for your answer.

28. What is special about the integral

$$\oint_C -y^3 dy + x^3 dx?$$

Give reasons for your answer.

29. **Area as a line integral** Show that if  $R$  is a region in the plane bounded by a piecewise-smooth simple closed curve  $C$ , then

$$\text{Area of } R = \oint_C x dy = - \oint_C y dx.$$

30. **Definite integral as a line integral** Suppose that a nonnegative function  $y = f(x)$  has a continuous first derivative on  $[a, b]$ . Let  $C$  be the boundary of the region in the  $xy$ -plane that is bounded below by the  $x$ -axis, above by the graph of  $f$ , and on the sides by the lines  $x = a$  and  $x = b$ . Show that

$$\int_a^b f(x) dx = - \oint_C y dx.$$

31. **Area and the centroid** Let  $A$  be the area and  $\bar{x}$  the  $x$ -coordinate of the centroid of a region  $R$  that is bounded by a piecewise-smooth simple closed curve  $C$  in the  $xy$ -plane. Show that

$$\frac{1}{2} \oint_C x^2 dy = - \oint_C xy dx = \frac{1}{3} \oint_C x^2 dy - xy dx = A\bar{x}.$$

32. **Moment of inertia** Let  $I_y$  be the moment of inertia about the  $y$ -axis of the region in Exercise 31. Show that

$$\frac{1}{3} \oint_C x^3 dy = - \oint_C x^2y dx = \frac{1}{4} \oint_C x^3 dy - x^2y dx = I_y.$$