

The Fundamental Theorem of Calculus, the normal form of Green's Theorem, and the Divergence Theorem all say that the integral of the differential operator  $\nabla \cdot$  operating on a field  $F$  over a region equals the sum of the normal field components over the boundary of the region. (Here we are interpreting the line integral in Green's Theorem and the surface integral in the Divergence Theorem as "sums" over the boundary.)

Stokes' Theorem and the tangential form of Green's Theorem say that, when things are properly oriented, the integral of the normal component of the curl operating on a field equals the sum of the tangential field components on the boundary of the surface.

The beauty of these interpretations is the observance of a single unifying principle which we might state as follows.

The integral of a differential operator acting on a field over a region equals the sum of the field components appropriate to the operator over the boundary of the region.

### Calculating Divergence

In Exercises 1–4, find the divergence of the field.

1. The spin field in Figure 16.14.
2. The radial field in Figure 16.13.
3. The gravitational field in Figure 16.9.
4. The velocity field in Figure 16.12.

### Using the Divergence Theorem to Calculate Outward Flux

In Exercises 5–16, use the Divergence Theorem to find the outward flux of  $F$  across the boundary of the region  $D$ .

5. **Cube**  $F = (y - x)\mathbf{i} + (z - y)\mathbf{j} + (y - x)\mathbf{k}$   
 $D$ : The cube bounded by the planes  $x = \pm 1, y = \pm 1$ , and  $z = \pm 1$
6.  $F = x^2\mathbf{i} + y^2\mathbf{j} + z^2\mathbf{k}$ 
  - a. **Cube**  $D$ : The cube cut from the first octant by the planes  $x = 1, y = 1$ , and  $z = 1$
  - b. **Cube**  $D$ : The cube bounded by the planes  $x = \pm 1, y = \pm 1$ , and  $z = \pm 1$
  - c. **Cylindrical can**  $D$ : The region cut from the solid cylinder  $x^2 + y^2 \leq 4$  by the planes  $z = 0$  and  $z = 1$
7. **Cylinder and paraboloid**  $F = y\mathbf{i} + xy\mathbf{j} - z\mathbf{k}$   
 $D$ : The region inside the solid cylinder  $x^2 + y^2 \leq 4$  between the plane  $z = 0$  and the paraboloid  $z = x^2 + y^2$
8. **Sphere**  $F = x^2\mathbf{i} + xz\mathbf{j} + 3z\mathbf{k}$   
 $D$ : The solid sphere  $x^2 + y^2 + z^2 \leq 4$
9. **Portion of sphere**  $F = x^2\mathbf{i} - 2xy\mathbf{j} + 3xz\mathbf{k}$   
 $D$ : The region cut from the first octant by the sphere  $x^2 + y^2 + z^2 = 4$
10. **Cylindrical can**  $F = (6x^2 + 2xy)\mathbf{i} + (2y + x^2z)\mathbf{j} + 4x^2y^3\mathbf{k}$   
 $D$ : The region cut from the first octant by the cylinder  $x^2 + y^2 = 4$  and the plane  $z = 3$
11. **Wedge**  $F = 2xz\mathbf{i} - xy\mathbf{j} - z^2\mathbf{k}$   
 $D$ : The wedge cut from the first octant by the plane  $y + z = 4$  and the elliptical cylinder  $4x^2 + y^2 = 16$
12. **Sphere**  $F = x^3\mathbf{i} + y^3\mathbf{j} + z^3\mathbf{k}$   
 $D$ : The solid sphere  $x^2 + y^2 + z^2 \leq a^2$
13. **Thick sphere**  $F = \sqrt{x^2 + y^2 + z^2}(x\mathbf{i} + y\mathbf{j} + z\mathbf{k})$   
 $D$ : The region  $1 \leq x^2 + y^2 + z^2 \leq 2$
14. **Thick sphere**  $F = (x\mathbf{i} + y\mathbf{j} + z\mathbf{k})/\sqrt{x^2 + y^2 + z^2}$   
 $D$ : The region  $1 \leq x^2 + y^2 + z^2 \leq 4$
15. **Thick sphere**  $F = (5x^3 + 12xy^2)\mathbf{i} + (y^3 + e^y \sin z)\mathbf{j} + (5z^3 + e^y \cos z)\mathbf{k}$   
 $D$ : The solid region between the spheres  $x^2 + y^2 + z^2 = 1$  and  $x^2 + y^2 + z^2 = 2$
16. **Thick cylinder**  $F = \ln(x^2 + y^2)\mathbf{i} - \left(\frac{2z}{x} \tan^{-1} \frac{y}{x}\right)\mathbf{j} + z\sqrt{x^2 + y^2}\mathbf{k}$   
 $D$ : The thick-walled cylinder  $1 \leq x^2 + y^2 \leq 2, -1 \leq z \leq 2$

### Properties of Curl and Divergence

17.  $\text{div}(\text{curl } G)$  is zero
- Show that if the necessary partial derivatives of the components of the field  $G = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$  are continuous, then  $\nabla \cdot \nabla \times G = 0$ .
  - What, if anything, can you conclude about the flux of the field  $\nabla \times G$  across a closed surface? Give reasons for your answer.
18. Let  $F_1$  and  $F_2$  be differentiable vector fields and let  $a$  and  $b$  be arbitrary real constants. Verify the following identities.
- $\nabla \cdot (aF_1 + bF_2) = a\nabla \cdot F_1 + b\nabla \cdot F_2$
  - $\nabla \times (aF_1 + bF_2) = a\nabla \times F_1 + b\nabla \times F_2$
  - $\nabla \cdot (F_1 \times F_2) = F_2 \cdot \nabla \times F_1 - F_1 \cdot \nabla \times F_2$
19. Let  $F$  be a differentiable vector field and let  $g(x, y, z)$  be a differentiable scalar function. Verify the following identities.
- $\nabla \cdot (gF) = g\nabla \cdot F + \nabla g \cdot F$
  - $\nabla \times (gF) = g\nabla \times F + \nabla g \times F$
20. If  $F = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$  is a differentiable vector field, we define the notation  $F \cdot \nabla$  to mean

$$M \frac{\partial}{\partial x} + N \frac{\partial}{\partial y} + P \frac{\partial}{\partial z}$$

For differentiable vector fields  $F_1$  and  $F_2$ , verify the following identities.

- $\nabla \times (F_1 \times F_2) = (F_2 \cdot \nabla)F_1 - (F_1 \cdot \nabla)F_2 + (\nabla \cdot F_2)F_1 - (\nabla \cdot F_1)F_2$
- $\nabla(F_1 \cdot F_2) = (F_1 \cdot \nabla)F_2 + (F_2 \cdot \nabla)F_1 + F_1 \times (\nabla \times F_2) + F_2 \times (\nabla \times F_1)$


### Theory and Examples

21. Let  $F$  be a field whose components have continuous first partial derivatives throughout a portion of space containing a region  $D$  bounded by a smooth closed surface  $S$ . If  $|F| \leq 1$ , can any bound be placed on the size of

$$\iiint_D \nabla \cdot F \, dV?$$

Give reasons for your answer.

22. The base of the closed cubelike surface shown here is the unit square in the  $xy$ -plane. The four sides lie in the planes  $x = 0$ ,  $x = 1$ ,  $y = 0$ , and  $y = 1$ . The top is an arbitrary smooth surface whose identity is unknown. Let  $F = x\mathbf{i} - 2y\mathbf{j} + (z + 3)\mathbf{k}$  and suppose the outward flux of  $F$  through side  $A$  is 1 and through side  $B$  is  $-3$ . Can you conclude anything about the outward flux through the top? Give reasons for your answer.



23. a. Show that the flux of the position vector field  $F = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  outward through a smooth closed surface  $S$  is three times the volume of the region enclosed by the surface.
- b. Let  $\mathbf{n}$  be the outward unit normal vector field on  $S$ . Show that it is not possible for  $F$  to be orthogonal to  $\mathbf{n}$  at every point of  $S$ .
24. **Maximum flux** Among all rectangular solids defined by the inequalities  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq 1$ , find the one for which the total flux of  $F = (-x^2 - 4xy)\mathbf{i} - 6yz\mathbf{j} + 12z\mathbf{k}$  outward through the six sides is greatest. What is the greatest flux?
25. **Volume of a solid region** Let  $F = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  and suppose that the surface  $S$  and region  $D$  satisfy the hypotheses of the Divergence Theorem. Show that the volume of  $D$  is given by the formula

$$\text{Volume of } D = \frac{1}{3} \iint_S F \cdot \mathbf{n} \, d\sigma$$

26. **Flux of a constant field** Show that the outward flux of a constant vector field  $F = C$  across any closed surface to which the Divergence Theorem applies is zero.
27. **Harmonic functions** A function  $f(x, y, z)$  is said to be *harmonic* in a region  $D$  in space if it satisfies the Laplace equation

$$\nabla^2 f = \nabla \cdot \nabla f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2} = 0$$

throughout  $D$ .

- Suppose that  $f$  is harmonic throughout a bounded region  $D$  enclosed by a smooth surface  $S$  and that  $\mathbf{n}$  is the chosen unit normal vector on  $S$ . Show that the integral over  $S$  of  $\nabla f \cdot \mathbf{n}$ , the derivative of  $f$  in the direction of  $\mathbf{n}$ , is zero.
- Show that if  $f$  is harmonic on  $D$ , then

$$\iint_S f \nabla f \cdot \mathbf{n} \, d\sigma = \iiint_D |\nabla f|^2 \, dV$$